arXiv:1110.2087v1 [math.DG] 10 Oct 2011

THE SPACETIME POSITIVE MASS THEOREM IN DIMENSIONS LESS THAN EIGHT

MICHAEL EICHMAIR, LAN-HSUAN HUANG, DAN A. LEE, AND RICHARD SCHOEN

Abstract. We prove the spacetime positive mass theorem in dimensions less than eight. This theorem states that for any asymptotically flat initial data set satisfying the dominant energy condition, the inequality $E \geq |P|$ holds, where (E, P) is the ADM energy-momentum vector of the initial data set. Previously, this theorem was proven only for spin manifolds [33]. Our proof is a modification of the minimal hypersurface technique that was used by the last author and S.-T. Yau to establish the time-symmetric case [25, 22]. Instead of minimal hypersurfaces, we use marginally outer trapped hypersurfaces (MOTS) whose existence is guaranteed by earlier work of the first author [12]. Since MOTS do not arise from a variational principle, an important part of our proof to introduce an appropriate substitute for the area functional used in the time-symmetric case. As part of our proof, we establish a density theorem of independent interest that allows us to reduce the general case of our theorem to the case of initial data that has harmonic asymptotics and satisfies the strict dominant energy condition. A refinement of the density argument allows us to approximate any given data set by one which is identical with a Kerr slice outside a compact set preserving the dominant energy condition. This enables us to give an alternative proof of the main theorem by reducing it to the positive energy theorem.

1. Introduction

The main result of this article is the following theorem.

Theorem 1 (Spacetime positive mass theorem)**.** *Let $3 \leq n < 8$, and let (M, g, k) be an n-dimensional asymptotically flat initial data set satisfying the dominant energy condition. Then*

$$E \geq |P|,$$

where (E, P) is the ADM energy-momentum vector of (M, g, k).

Date: October 11, 2011.

The first author acknowledges the support of NSF grant DMS-0906038. The second author acknowledges the support of NSF grant DMS-1005560. The third author acknowledges the support of NSF grant DMS-0903467. The fourth author acknowledges the support of NSF grant DMS-0604960.

Precise definitions of all technical terms used in this article are given in Section 2.

We briefly survey earlier results: A particularly important special case of this theorem is when $k \equiv 0$, which is called the time-symmetric case, or sometimes the Riemannian case. In this case, we have $P = 0$, and the dominant energy condition reduces to the assumption of nonnegative scalar curvature of g. The last author and S.-T. Yau proved the time-symmetric case of the theorem in dimension three in two articles from 1979 and 1981 [25, 26]. In [28] they generalized their proof of the time-symmetric case to dimensions less than 8 without any topological assumption, as explained in detail in [22]. In 1981, they considered the general case $k \not\equiv 0$ in dimension three and succeeded in proving that $E \geq 0$ by solving Jang's equation [27]. Later, E. Witten discovered a completely different proof that $E \geq |P|$ in dimension three [33]. Witten's technique generalizes to all higher dimensions, as long as the manifold is *spin* [5, 9]. In higher dimensions, a complication arises in the Schoen-Yau argument due to possible singularities of minimal hypersurfaces. Two different approaches for handling this complication have been announced by J. Lohkamp in a 2006 preprint [19] and by the last author in 2009. The first author generalized the spacetime $E \geq 0$ theorem to dimensions less than 8 (without spin assumption) in his thesis (see [11]).

For earlier history of this problem, see [25]. The $E \geq 0$ theorem is sometimes referred to as the "positive mass theorem," but we prefer to refer to it more accurately as the "positive energy theorem," and reserve the phrase "positive mass theorem" for the $E \geq |P|$ theorem, which could also reasonably be called the "future timelike energy-momentum theorem." One can see that there is a natural relationship between these two theorems by "boosting" an initial data slice of a spacetime. In the case of vacuum initial data this approach can be made rigorous by using [8] to approximate the data by an initial data set which coincides with a Kerr data set outside a compact set. The boost can then be done near infinity in the Kerr spacetime. In the last section of the paper we show that this argument can be generalized to the case in which the data satisfies the dominant energy condition. More precisely we show that a general initial data set satisfying the dominant energy condition can be approximatied by one which is identical with a Kerr slice near infinity and still satisfies the dominant energy condition. This gives an alternative proof of the positive mass theorem from the positive energy theorem. It is discussed in the final section of the paper. We emphasize that the proof presented in this paper is self-contained and gives a new proof of the positive energy theorem as well.

We point out that our main theorem does not include a rigidity statement. The expected rigidity theorem, which has been proven in the spin case [34, 6, 7], states that if, in addition to the hypotheses in the main theorem, we have $E = |P|$, then $E = |P| = 0$ and (M, g) isometrically embeds into Minkowski space with k as its second fundamental form.

We briefly review the minimal hypersurface proof of the time-symmetric positive mass theorem in [25, 22], since our proof is closely related. The argument is by induction on the dimension $3 \leq n < 8$, and proceeds by contradiction. Suppose that there exists an asymptotically flat Riemannian manifold (M, g) with nonnegative scalar curvature $R_g \geq 0$ and negative mass $E < 0$. By a density argument [26], one may assume that (M, g) has *harmonic asymptotics* and positive scalar curvature. The harmonic asymptotics and $E < 0$ assumptions imply that for large enough Λ, the coordinate planes $x^n = \pm\Lambda$ are barriers for minimal hypersurfaces. Consider an $(n-1)$-dimensional vertical cylinder ∂C_ρ of large radius ρ in the asymptotically flat coordinates. For each $h \in [-\Lambda, \Lambda]$, let $\Sigma_{\rho,h}$ be an area-minimizing hypersurface whose boundary is equal to the height h sphere on ∂C_ρ. As long as $n < 8$, this area-minimizing hypersurface must be smooth. Each $\Sigma_{\rho,h}$ must lie between the barrier planes $x^n = \pm\Lambda$, and the area $\mathcal{H}^{n-1}(\Sigma_{\rho,h})$ is actually minimized over h by some $h_\rho \in (-\Lambda, \Lambda)$. This means that this Σ_{ρ,h_ρ} has the property that

$$\left.\frac{d^2}{dt^2}\right|_{t=0} \mathcal{H}^{n-1}(\Phi_t(\Sigma_{\rho,h_\rho})) \geq 0, \tag{1}$$

for any variation Φ_t of Σ_{ρ,h_ρ} that is equal to vertical translation along ∂C_ρ. One can then extract a smooth subsequential limit Σ_∞ of Σ_{ρ,h_ρ} as $\rho \to \infty$. This Σ_∞ is itself an $(n-1)$-dimensional asymptotically flat manifold with energy equal to zero. Moreover, Σ_∞ is a stable minimal hypersurface of M. Not only does (1) imply that it is stable with respect to compact variations, but it is also stable with respect to variations that are (sufficiently close to) vertical translations outside a compact set. Using the (by now) well-established relationship between the stability of minimal hypersurfaces and scalar curvature, the stability of Σ_∞ allows one to construct a conformal factor that changes the metric on Σ_∞ to one with zero scalar curvature. The stability with respect to variations that are close to vertical translations is then used to show that the conformal factor must *decrease* the energy of Σ_∞, thereby violating the time-symmetric positive mass theorem in dimension $n-1$. For the base case of the induction, when $n = 3$, one can show that the stability of Σ_∞ combined with its asymptotics at infinity are incompatible with the Gauss-Bonnet Theorem. Note that when $n = 3$, choosing a special height h_ρ turns out to be unnecessary.

We now compare this with our approach to the spacetime positive mass theorem, which is essentially a *generalization* of the proof described above. In particular, it does not use the time-symmetric positive mass theorem as an input, as was done in the Jang equation approach of [27]. Again, the proof is by contradiction. Let (M^n, g, k) be an asymptotically flat initial data set satisfying the dominant energy condition and $E < |P|$. By a density argument, which we give in Section 6, we may assume that (M, g, k) has *harmonic asymptotics* and satisfies the strict dominant energy condition $\mu > |J|$. We may assume further that P points in the vertical direction $-\partial_n$ in the asymptotic coordinate chart. The harmonic asymptotics and $E < |P|$ assumptions imply that for large enough Λ, the coordinate planes $x^n = \pm\Lambda$ are barriers for *marginally outer trapped hypersurfaces* (MOTS). Again, we consider an $(n-1)$-dimensional vertical cylinder ∂C_ρ of large radius ρ in the asymptotically flat coordinates. For each $h \in [-\Lambda, \Lambda]$, the existence theory of [12] allows us to use the barriers to construct a MOTS $\Sigma_{\rho,h}$ whose boundary is equal to the height h sphere on ∂C_ρ. As long as $n < 8$, this MOTS is smooth, because it has a λ-minimizing property (see Section 2). Moreover, $\Sigma_{\rho,h}$ lies between the planes $x^n = \pm\Lambda$ and is *stable* in the sense of MOTS with boundary [16] (see Section 2). Since MOTS are not known to arise from a variational principle in general, we cannot find a suitable height h_ρ simply by minimizing some quantity. To this end, we introduce a new functional $\mathcal{F}$ on hypersurfaces with boundary on ∂C_ρ, for which we can find $h_\rho \in (-\Lambda, \Lambda)$ with (roughly) the property that

$$\left.\frac{d}{dh}\right|_{h=h_\rho} \mathcal{F}(\Sigma_{\rho,h}) \geq 0. \tag{2}$$

Inequality (2) in conjunction with the MOTS-stability of $\Sigma_{\rho,h}$ then plays a role similar to that of (1) in the time-symmetric case. Note that the h_ρ selected in the time-symmetric case by minimization would satisfy (2) in our more general argument. As before, we extract a smooth subsequential limit Σ_∞ of Σ_{ρ,h_ρ} as $\rho \to \infty$. This Σ_∞ is itself an $(n-1)$-dimensional asymptotically flat manifold with energy equal to zero, and Σ_∞ is a stable MOTS in M. Using the relationship between stability of MOTS and scalar curvature established in [17], one can construct a conformal factor that changes the metric on Σ_∞ to one with zero scalar curvature. Finally, (2) plays the role of (1) in establishing that the conformal factor must decrease the energy of Σ_∞, thereby violating the *time-symmetric* positive mass theorem in dimension $n-1$.

As in the time-symmetric case, the delicate height-picking argument is not required when $n = 3$. As before, we use the asymptotics of Σ_∞ and MOTS-stability to produce a contradiction to the Gauss-Bonnet theorem.

The structure of the paper is as follows. Section 2 sets up the basic definitions and recalls some useful background material. Section 3 establishes the existence of the MOTS needed for the proof. Section 4 completes the $n = 3$ case of the main theorem (Theorem 1). The basic $n = 3$ argument, which is explained in detail in Sections 3 and 4, was first sketched out in [23, Section 7.2]. Section 5 contains the parts of the proof that are specific to dimensions higher than 3, including the height-picking procedure alluded to above. Finally, the last section contains the relevant density theorem and its proof, which may be of independent interest.

2. Definitions, notation, and basic facts

In this section we review some of the terminology used in this article, as well as some basic facts.

Definition 1. Let B be a closed ball in $\mathbb{R}^n$ centered at the origin, and consider the exterior region $\mathbb{R}^n \smallsetminus B$. For any $k \in \mathbb{N}$, $p \geq 1$, $q \in \mathbb{R}$, we define the *weighted Sobolev space* $W^{k,p}_{-q}(\mathbb{R}^n \smallsetminus B)$ as the space of Lebesgue measurable functions f on $\mathbb{R}^n \smallsetminus B$ such that

$$\|f\|_{W^{k,p}_{-q}(\mathbb{R}^n \smallsetminus B)} := \left(\int_{\mathbb{R}^n \smallsetminus B} \sum_{|\alpha| \leq k} \left(|\partial^\alpha f(x)| |x|^{|\alpha|+q} \right)^p |x|^{-n} \, dx \right)^{\frac{1}{p}} < \infty.$$

Now suppose that M is a $\mathcal{C}^k$ manifold such that there is a compact set $K \subset M$ and a diffeomorphism $M \smallsetminus K \cong \mathbb{R}^n \smallsetminus B$. Then one can easily define $W^{k,p}_{-q}(M)$ in a routine manner, by choosing an atlas for M that consists of the diffeomorphism $M \smallsetminus K \cong \mathbb{R}^n \smallsetminus B$ together with finitely many precompact charts, and then using the $W^{k,p}_{-q}(\mathbb{R}^n \smallsetminus B)$ norm on the noncompact chart while using the $W^{k,p}$ norm on the other ones. Note that the space $W^{k,p}_{-q}(M)$ and its topology depend only on the diffeomorphism $M \smallsetminus K \cong \mathbb{R}^n \smallsetminus B$. The definition of $W^{k,p}_{-q}(M)$ also extends to tensors simply by considering their components with respect to these coordinate charts. We will usually write $W^{k,p}_{-q}$ for $W^{k,p}_{-q}(M)$ when the context is clear.

Definition 2. Consider the exterior region $\mathbb{R}^n \smallsetminus B$. For any $k \in \mathbb{N}$, $\alpha \in (0,1)$, $q \in \mathbb{R}$, we define the *weighted Hölder space* $\mathcal{C}^{k,\alpha}_{-q}(\mathbb{R}^n \smallsetminus B)$ to be the space of functions f on $\mathbb{R}^n \smallsetminus B$ such that

$$\|f\|_{\mathcal{C}^{k,\alpha}_{-q}(\mathbb{R}^n \smallsetminus B)} := \sum_{|\alpha| \leq k} \sup_x \left| |x|^{|\alpha|+q} \partial^\alpha f(x) \right| + \sum_{|\alpha|=k} \left[|x|^{|\alpha|+q} \partial^\alpha f(x) \right]_\alpha < \infty.$$

If M is a $\mathcal{C}^{k,\alpha}$ manifold such that there is a compact set K and a diffeomorphism $M \smallsetminus K \cong \mathbb{R}^n \smallsetminus B$, then one can define $\mathcal{C}^{k,\alpha}_{-q}(M)$ just as we defined $W^{k,p}_{-q}(M)$ above.

Definition 3. Let $n \geq 3$ and $\alpha \in (0,1)$. An *initial data set* is a complete $\mathcal{C}^{3,\alpha}$ manifold M^n equipped with a $\mathcal{C}^{2,\alpha}_{loc}$ Riemannian metric g and a $\mathcal{C}^{1,\alpha}_{loc}$ symmetric $(0,2)$-tensor k. On an initial data set, one can define the *mass density* μ and the *current density* J by

$$\begin{aligned} \mu &= \tfrac{1}{2} \left(R_g - |k|^2_g + (\operatorname{tr}_g k)^2 \right) \\ J &= \operatorname{div}_g k - d(\operatorname{tr}_g k). \end{aligned}$$

We say that (M, g, k) satisfies the *dominant energy condition* if

$$\mu \geq |J|_g.$$

It is often convenient to consider the *momentum tensor*

$$\pi = k - (\operatorname{tr}_g k)g.$$

Note that π contains the same information as k since $k = \pi - \frac{1}{n-1}(\operatorname{tr}_g \pi)g$.

Let $p > n$, $q > \frac{n-2}{2}$, and $q_0 > 0$. We say that the initial data set (M, g, k) is *asymptotically flat*[1] if there is a compact set $K \subset M$ and a $\mathcal{C}^{3,\alpha}$ diffeomorphism $M \smallsetminus K \cong \mathbb{R}^n \smallsetminus B$ for some closed ball $B \subset \mathbb{R}^n$, such that in these coordinates, for $i, j = 1, 2, \ldots, n$,

$$(g_{ij} - \delta_{ij}, k_{ij}) \in W^{2,p}_{-q}(\mathbb{R}^n \smallsetminus B) \times W^{1,p}_{-1-q}(\mathbb{R}^n \smallsetminus B)$$

and

$$(\mu, J) \in \mathcal{C}^{0,\alpha}_{-n-q_0}.$$

We sometimes say that (M, g, k) is asymptotically flat *of type* (p, q, q_0, α).

Under these hypotheses, one can define the *ADM energy* E and *ADM momentum* P to be

$$E = \tfrac{1}{2(n-1)\omega_{n-1}} \lim_{r\to\infty} \int_{|x|=r} \sum_{i,j=1}^{n} (g_{ij,i} - g_{ii,j})\nu_0^j \, d\mathcal{H}_0^{n-1}$$

$$P_i = \tfrac{1}{(n-1)\omega_{n-1}} \lim_{r\to\infty} \int_{|x|=r} \sum_{j=1}^{n} \pi_{ij}\nu_0^j \, d\mathcal{H}_0^{n-1}, \qquad i = 1, 2, \ldots, n$$

where $\nu_0^j = \frac{x^j}{|x|}$ is the Euclidean outward unit normal, the measure $\mathcal{H}_0^{n-1}$ is Euclidean Hausdorff measure, and ω_{n-1} is the volume of the standard unit sphere in $\mathbb{R}^n$.

Remark. Theorem 1 still holds if we allow multiple asymptotically flat ends in the definition of initial data sets. We simply use the large celestial spheres in the "other" ends as barriers in the proof of Lemma 6.

In order to execute our main argument, we require better asymptotic behavior.

Definition 4. Let $n \geq 3$, and let (M^n, g, k) be an asymptotically flat initial data set of type (p, q, q_0, α). We say that (M, g, k) has *harmonic asymptotics* if there exists a $\mathcal{C}^{3,\alpha}$ diffeomorphism as in the definition of asymptotic flatness, as well as a $\mathcal{C}^{2,\alpha}_{2-n}$ function u and a $\mathcal{C}^{2,\alpha}_{2-n}$ vector field Y such that for $i, j = 1, 2, \ldots, n$,

$$\begin{aligned} u(x) &= 1 + a|x|^{2-n} + O^{2,\alpha}(|x|^{1-n}) \\ Y_i(x) &= b_i|x|^{2-n} + O^{2,\alpha}(|x|^{1-n}) \\ g_{ij} &= u^{\frac{4}{n-2}}\delta_{ij} \\ \pi_{ij} &= u^{\frac{2}{n-2}}\left[(L_Y\delta)_{ij} - (\operatorname{div}_\delta Y)\delta_{ij}\right] \end{aligned}$$

---

[1] Note that the term *asymptotically flat* has various incompatible definitions in the literature.

where a, b_i are constants, and L_Y is the Lie derivative. The $O^{2,\alpha}(|x|^{1-n})$ indicates that the function lies in the weighted Hölder space $\mathcal{C}^{2,\alpha}_{1-n}$.

Notation. Let (M^n, g, k) be an initial data set, and let Σ^{n-1} be a $\mathcal{C}^{3,\alpha}$ hypersurface with boundary in M^n, equipped with a global unit normal ν. Let D denote the covariant derivative with respect to the ambient metric on M. (In contrast, we will usually use ∇ when differentiating functions.) We define the second fundamental form B_Σ and shape operator S_Σ of Σ using the convention

$$B_\Sigma(X, Y) = \langle S_\Sigma(X), Y\rangle = \langle D_X \nu, Y\rangle$$

for vector fields X, Y tangent to Σ, where the angle brackets denote the inner product g. We define the mean curvature scalar H_Σ to be the trace of S_Σ. Using this convention, if ν is the upward-pointing normal of the upper hemisphere in $\mathbb{R}^n$, the mean curvature is positive. We also define the *expansion*

$$\theta^+_\Sigma = H_\Sigma + \operatorname{tr}_\Sigma k,$$

where $\operatorname{tr}_\Sigma k$ denotes the trace over the tangent space of Σ. If θ^+_Σ vanishes on all of Σ, then we say that Σ is a *marginally outer trapped hypersurface*, or *MOTS* for short.[2] Note that the property of being a MOTS depends on the choice of normal. For any vector field X defined along Σ but not necessarily tangent to it, we define

$$\operatorname{div}_\Sigma X = \sum_{i=1}^{n-1} \langle D_{e_i} X, e_i\rangle,$$

where $e_1, \ldots, e_{n-1}$ is a local orthonormal frame of Σ.

Notation. Given a vector field X defined along Σ, we can decompose X into its normal and tangential components

$$X = \varphi\nu + \hat{X}.$$

Throughout this paper, whenever there is a vector field with the variable name X on a hypersurface Σ, the function φ and the vector field $\hat{X}$ are defined according to this relationship. We use η to denote the outward unit normal of $\partial\Sigma$ in Σ.

The expression for the linearization of the expansion stated in the following proposition generalizes the well-known variation formula for the mean curvature. See, for example, [17].

Proposition 2. *Let Σ^{n-1} be a two-sided hypersurface with boundary in an initial data set (M^n, g, k), and let ν be a continuous unit normal field along Σ. Let $X \in \Gamma(TM)$ be a $\mathcal{C}^2$ vector field, and let Φ_t be the flow generated by X. We can compute the*

---

[2] While both the nomenclature and the acronym are slightly inappropriate in our context, we find it convenient to use these terms.

expansion $\theta^+_{\Sigma_t}$ *of the push forward* $\Sigma_t := \Phi_t(\Sigma)$ *with respect to the unit normal that points in the direction of* $\Phi_{t*}(\nu)$ *and pull it back to a function on* Σ *using* Φ_t*. The derivative of this function in* t *at* $t = 0$ *is denoted by* $D\theta^+|_\Sigma(X)$ *and one has that*

$$D\theta^+|_\Sigma(X) = -\Delta_\Sigma\varphi + 2\langle W_\Sigma, \nabla\varphi\rangle + (\operatorname{div}_\Sigma W_\Sigma - |W_\Sigma|^2 + Q_\Sigma)\varphi + \nabla_{\hat{X}}\theta^+_\Sigma, \quad (3)$$

where

$$Q_\Sigma = \tfrac{1}{2}R_\Sigma - \mu - J(\nu) - \tfrac{1}{2}|k_\Sigma + B_\Sigma|^2.$$

Here, k_Σ *denotes the restriction of* k *to vectors tangent to* Σ*, and* W_Σ *is the tangential vector field on* Σ *that is dual to the* 1*-form* $k(\nu, \cdot)$ *along* Σ*.*

Notation. Throughout this paper, we will frequently drop the Σ subscripts when there is minimal ambiguity. Everything is computed with respect to the metric g unless stated otherwise, so we will not write g explicitly in any notation that we use. In particular, we use $\mathcal{H}$ to denote Hausdorff measure with respect to the metric g, unless indicated otherwise.

Definition 5. Let Σ be a MOTS in an initial data set (M, g, k). We define the operator

$$L_\Sigma v := -\Delta_\Sigma v + 2\langle W_\Sigma, \nabla v\rangle + (\operatorname{div}_\Sigma W_\Sigma - |W_\Sigma|^2 + Q_\Sigma)v, \quad (4)$$

where v is a function on Σ. Although this operator is not self-adjoint, the Krein-Rutman Theorem shows that there exists a (Dirichlet) eigenvalue with minimal real part, which is called the *principal (Dirichlet) eigenvalue* of L_Σ. Furthermore, this eigenvalue is real, and if Σ is connected, the corresponding eigenspace is a one-dimensional space generated by a $\mathcal{C}^{2,\alpha}$ *principal eigenfunction* that is positive on the interior of Σ [16, p. 3]. If the principal eigenvalue is nonnegative, we say that Σ is a *stable* MOTS. For a closed MOTS, this concept goes back to [16, 2]. It is easy to see that this generalizes the notion of stability of minimal hypersurfaces with boundary.

Proposition 3. *Let* Σ *be a stable MOTS in an initial data set* (M, g, k)*. For any compactly supported* $\mathcal{C}^1$ *function* v *on* Σ *that vanishes along* $\partial\Sigma$*, we have that*

$$\int_\Sigma (|\nabla v|^2 + Q_\Sigma v^2)\, d\mathcal{H}^{n-1} \geq 0. \quad (5)$$

This result follows from an argument in [17], *cf.* the proof of Lemma 15 below.

We state the following geometric variant of the strong maximum principle for ordered hypersurfaces that are subsolutions and supersolutions of the same prescribed mean curvature equation. We refer to [21, Lemma 1], [3, Proposition 3.1], and [4, Proposition 2.4] for similar results and partial proofs. It is important to pay attention to the choice of normal here, and a good example to have in mind is the following: Consider a sphere of radius 2 that is tangent to a sphere of radius 1 in Euclidean space. Either the smaller one is enclosed by the larger one or lies outside it. The (obvious) conclusion from the lemma in this simple example is that the larger sphere cannot lie inside the smaller one.

Proposition 4 (strong maximum principle)**.** *Let g be a complete $\mathcal{C}^2$ Riemannian metric on $M := B_2^{n-1}(0) \times (-2,2) \subset \mathbb{R}^n$, let F be a $\mathcal{C}^1$ function on the unit sphere bundle $\mathbb{S}_g^{n-1}(TM)$, and let $u_1, u_2 \in \mathcal{C}^2(\bar{B}_1^{n-1}(0))$ be such that $-1 \le u_1(x') \le u_2(x') \le 1$ for all $x' \in \bar{B}^{n-1}(0,1)$. We assume that the hypersurfaces with boundary $\Sigma_i :=$ graph$(u_i) \subset M$ are such that $H_{\Sigma_1}(x) \le F(x, \nu_{\Sigma_1}(x))$ for all $x \in \Sigma_1$ and $H_{\Sigma_2}(x) \ge F(x, \nu_{\Sigma_2}(x))$ for all $x \in \Sigma_2$. Here, the mean curvature is computed using the upward pointing unit normals. If Σ_1 and Σ_2 intersect at an interior point or are tangent to each other at a boundary point, then they must be equal.*

Remark. We will use this proposition in the following context: We will have two open sets $\Omega_1 \subset \Omega_2$ such that Σ_1 and Σ_2 lie on $\partial\Omega_1$ and $\partial\Omega_2$, respectively, and $\theta^+_{\Sigma_1} \le 0$ while $\theta^+_{\Sigma_2} \ge 0$.

Next, we recall a few facts about conformal changes of the metric. In general, given two metrics on M related by $g_2 = u^{\frac{4}{n-2}} g_1$, the scalar curvatures are related by the formula

$$R_2 = u^{-\frac{n+2}{n-2}} \left(R_1 u - \tfrac{4(n-1)}{n-2} \Delta_1 u \right). \tag{6}$$

Given any hypersurface Σ in M, its unit normals satisfy $\nu_2 = u^{\frac{-2}{n-2}} \nu_1$, while the mean curvatures are related by

$$H_2 = u^{\frac{-2}{n-2}} \left(H_1 + \tfrac{2(n-1)}{n-2} u^{-1} \nabla_{\nu_1} u \right). \tag{7}$$

Also, if (M^n, g_1) is an asymptotically flat Riemannian manifold and $u = 1 + a|x|^{2-n} + O^{2,\alpha}(|x|^{1-n})$ is $\mathcal{C}^{2,\alpha}$, then (M^n, g_2) is also asymptotically flat, and the energies are related by the formula

$$\begin{aligned} E_2 &= E_1 - \tfrac{2}{(n-2)\omega_{n-1}} \lim_{r\to\infty} \int_{|x|=r} u \nabla_{\nu_1} u \, d\mathcal{H}_1^{n-1} && (8) \\ &= E_1 + 2a. && (9) \end{aligned}$$

Finally, the proof of the positive mass theorem in the time-symmetric case uses regularity and compactness properties of minimizing hypersurfaces. Since MOTS are not known to obey a useful variational principle, we rely on the theory of almost minimizing currents:

Definition 6 ([10])**.** Let (M^n, g) be a complete Riemannian manifold, and let T be an integral k-current in M. Let $U \subset M$ be an open set such that $\mathrm{spt}(\partial T) \cap U = \emptyset$. Then T is *λ-minimizing in U* if, for every integral $(k+1)$-current X with support in U, we have

$$\mathbf{M}_U(T) \le \mathbf{M}_U(T + \partial X) + \lambda \mathbf{M}_U(X).$$

Here, $\mathbf{M}_U$ denotes the mass of a current in U.

This almost minimizing property was introduced and studied systematically in [10]. In [12] it was shown that the λ-minimizing property is a natural feature of the MOTS that arise in the existence theory developed there, despite the absence of a useful variational characterization. The elementary properties of λ-minimizing boundaries that are needed in this paper are also summarized in [12, Appendix A] with further precise references to the geometric measure theory literature.

3. Construction of MOTS

Our proof of Theorem 1 will be by induction on dimension and contradiction. Let $3 \le n < 8$, and suppose there exists an asymptotically flat initial data set (M^n, g, k) of type (p, q, q_0, α) satisfying the dominant energy condition, but $E < |P|$. For the case $n = 3$, we will obtain a contradiction to the Gauss-Bonnet Theorem in Section 4, and for $3 < n < 8$, we will obtain a contradiction to the time-symmetric case of Theorem 1 in dimension $n-1$. By the density theorem (Theorem 18), we can assume without loss of generality, that (g, k) has harmonic asymptotics and satisfies the strict dominant energy condition $\mu > |J|_g$. Specifically, we can choose asymptotically flat coordinates $(x^1, \dots, x^n)$ on $M \smallsetminus K \cong_x \mathbb{R}^n \smallsetminus B$ for some closed ball B such that on $\mathbb{R}^n \smallsetminus B$, we have, for $i, j = 1, 2, \dots, n$,

$$\begin{aligned} g_{ij} &= u^{\frac{4}{n-2}} \delta_{ij} \\ \pi_{ij} &= u^{\frac{2}{n-2}} \left[(L_Y \delta)_{ij} - (\operatorname{div}_\delta Y) \delta_{ij} \right] \end{aligned}$$

for some $u, Y \in \mathcal{C}^{2,\alpha}_{2-n}$ satisfying

$$\begin{aligned} u(x) &= 1 + a|x|^{2-n} + O^{2,\alpha}(|x|^{1-n}) \\ Y_i(x) &= b_i |x|^{2-n} + O^{2,\alpha}(|x|^{1-n}). \end{aligned}$$

Without loss of generality, we assume that $P = (0, \dots, 0, -|P|)$.

3.1. **Existence of horizontal barriers.**

Lemma 5. *Let (M^n, g, k) be as described above. Then, for sufficiently large Λ, we have that $\theta^+_{\{x^n = \Lambda\}} > 0$ and $\theta^+_{\{x^n = -\Lambda\}} < 0$ where the expansion is computed with respect to the upward pointing unit normal.*

Proof. It follows from (9) that the $|x|^{2-n}$ coefficient of the function u is just $a = \frac{E}{2}$. We claim that the $|x|^{2-n}$ coefficient of Y_i is $b_i = -\frac{n-1}{n-2} P_i$, i.e. that $b_n = \frac{n-1}{n-2}|P|$ and

$b_i = 0$ for $i < n$. To see this, note that

$$\begin{aligned}(n-1)\omega_{n-1}P_i &= \lim_{r\to\infty}\int_{|x|=r} \pi_{ij}\nu_0^j\, d\mathcal{H}_0^{n-1}\\ &= \lim_{r\to\infty}\int_{|x|=r} (Y_{i,j}+Y_{j,i}-(\operatorname{div}_\delta Y)\delta_{ij})\,\nu_0^j\, d\mathcal{H}_0^{n-1}\\ &= \lim_{r\to\infty}\int_{|x|=r} (2-n)|x|^{1-n}\left[b_i(\nu_0)_j + b_j(\nu_0)_i - b_k\nu_0^k\delta_{ij} + O(|x|^{-1})\right]\nu_0^j\, d\mathcal{H}_0^{n-1}\\ &= -(n-2)b_i\omega_{n-1}.\end{aligned}$$

We claim that

$$\theta^+_{\{x^n=\Lambda\}} = H_{\{x^n=\Lambda\}} + \operatorname{tr}_{\{x^n=\Lambda\}}(k) = (n-1)(|P|-E)\Lambda|x|^{-n} + O(|x|^{-n}). \tag{10}$$

To see this, we use harmonic asymptotics and formula (7) to compute

$$\begin{aligned}H_{\{x^n=\Lambda\}} &= \tfrac{2(n-1)}{n-2}u^{\frac{-2}{n-2}-1}\partial_n u\\ &= \tfrac{2(n-1)}{n-2}u^{\frac{-2}{n-2}-1}[(2-n)a|x|^{-n}x^n + O(|x|^{-n})]\\ &= -2(n-1)a|x|^{-n}x^n + O(|x|^{-n})\\ &= -(n-1)E|x|^{-n}\Lambda + O(|x|^{-n}).\end{aligned}$$

To compute $\operatorname{tr}_{\{x^n=\Lambda\}}(k)$, first note that

$$k_{ij} = u^{\frac{2}{n-2}}\left[(L_Y\delta)_{ij} - \tfrac{1}{n-1}(\operatorname{div}_\delta Y)\delta_{ij}\right].$$

So we have

$$\begin{aligned}\operatorname{tr}_{\{x^n=\Lambda\}}(k) &= \sum_{i,j=1}^{n-1} g^{ij}k_{ij}\\ &= \sum_{i,j=1}^{n-1} u^{\frac{-2}{n-2}}\delta^{ij}[Y_{i,j}+Y_{j,i} - \tfrac{1}{n-1}(\operatorname{div}_\delta Y)\delta_{ij}]\\ &= \sum_{i,j=1}^{n-1} u^{\frac{-2}{n-2}}\delta^{ij}\left[\tfrac{-1}{n-1}Y_{n,n}\delta_{ij} + O(|x|^{-n})\right] \text{ since } Y_i = O^1(|x|^{1-n}) \text{ for } i<n\\ &= -Y_{n,n} + O(|x|^{-n})\\ &= (n-2)b_n|x|^{-n}x^n + O(|x|^{-n})\\ &= (n-1)|P||x|^{-n}\Lambda + O(|x|^{-n}),\end{aligned}$$

completing the proof of (10). Note that (10) shows that that for large enough Λ one has $\theta^+_{\{x^n=\Lambda\}} > 0$. The proof that $\theta^+_{\{x^n=-\Lambda\}} < 0$ is similar. □



FIGURE 1. The expansion θ^+ is computed with respect to the indicated unit normals.

3.2. **Existence of MOTS.**

Notation. We now fix Λ large enough so that Lemma 5 applies. For large ρ, we define $C_\rho := K \cup x^{-1}\{(x^1, \ldots, x^{n-1}, x^n) \,|\, (x^1)^2 + \ldots + (x^{n-1})^2 < \rho^2\}$ to be the region horizontally bounded by a vertical coordinate cylinder of radius ρ, and we define $C_{\rho,h}$ to be the part of C_ρ lying between the planes $x^n = \pm h$. Define $\Gamma_{\rho,h} := \partial C_\rho \cap \{x^n = h\}$.

Lemma 6. *Let (M^n, g, k) and Λ be as described above. For every sufficiently large ρ and all $h \in [-\Lambda, \Lambda]$ there exists a $\mathcal{C}^{3,\alpha}$ stable MOTS $\Sigma_{\rho,h}$ in $C_{\rho,\Lambda}$ whose boundary equals $\Gamma_{\rho,h}$ and which meets ∂C_ρ transversely. Each $\Sigma_{\rho,h}$ is a λ-minimizing (relative) boundary in $C_{\rho,2\Lambda}$ where λ depends only on $|k|_{\mathcal{C}^0}$. Moreover, we can choose $\{\Sigma_{\rho,h}\}_{|h|\le\Lambda}$ so that if $-\Lambda \le h_1 < h_2 \le \Lambda$, then Σ_{ρ,h_1} lies strictly "below" Σ_{ρ,h_2} in the sense of relative boundaries in $C_{\rho,\Lambda}$.*

Remark. The regularity (and later, convergence) of the MOTS $\Sigma_{\rho,h}$ is the only place in the proof of Theorem 1 where the assumption $n < 8$ is used. For $n > 8$, the lemma still holds except that the λ-minimizing boundaries $\Sigma_{\rho,h}$ are only regular away from a thin singular set.

Proof. First observe that by the decay conditions on g and k, the coordinate cylinder ∂C_ρ has $\theta^+ > 0$ with respect to the outward normal and $\theta^+ < 0$ with respect to the inward normal.

We would like to solve the MOTS Plateau problem with boundary equal to $\Gamma_{\rho,h}$ for each $h \in [-\Lambda, \Lambda]$. Note that $\Gamma_{\rho,h}$ divides $\partial C_{\rho,2\Lambda}$ into a top piece and a bottom piece.

According to [13] the MOTS Plateau problem is solvable if the top piece has $\theta^+ > 0$ with respect to the outward normal of $\partial C_{\rho,2\Lambda}$ and the bottom piece has $\theta^+ < 0$ with respect to the inward normal of $\partial C_{\rho,2\Lambda}$. By the observation above and Lemma 5, the two pieces of $\partial C_{\rho,2\Lambda}$ satisfy the desired trapping conditions, with the exception of the corners where ∂C_ρ intersects $\{x^n = \pm 2\Lambda\}$. See Figure 1. Intuitively, the corners do not cause a problem because the (singular) distributional mean curvature there has a favorable sign. Therefore one could "round-off" the corners as in Figure 1. Alternatively, we observe that the proof in [12] can easily accommodate the corners: Just as in that proof, we use the trapping of the cylindrical and horizontal pieces of $\partial C_{\rho,2\Lambda}$ to construct barriers that have appropriate blow-up behavior for Jang's equation, and then we simply combine these barriers as in [13, Lemma 4.1]. Specifically, there exists a $\mathcal{C}^{3,\alpha}$ family of MOTS $\{\Sigma_{\rho,h}\}_{|h|\leq\Lambda}$ with $\partial\Sigma_{\rho,h} = \Gamma_{\rho,h}$ such that each $\Sigma_{\rho,h}$ is a λ-minimizing relative boundary in $C_{\rho,2\Lambda}$, where $\lambda = \lambda(|k|_{\mathcal{C}^0})$.

Moreover, it can be seen from the construction in [12] that the regions $\Omega_{\rho,h} \subset C_{\rho,2\Lambda}$ bounded by the $\Sigma_{\rho,h}$'s can be taken to be ordered, so that $\Omega_{\rho,h_1} \subset \Omega_{\rho,h_2}$ whenever $-\Lambda \leq h_1 \leq h_2 \leq \Lambda$. To see this, note that the supersolutions $\overline{u}_{t,\rho,h}$ used in the construction of solutions to the regularized Jang's equation that lead to blow up in [12, Lemma 4.1, bottom of p. 570] can be taken to be pointwise decreasing in the parameter h, so that the corresponding Perron solutions $\overline{u}^P_{t,\rho,h}$ are decreasing in h and hence their epigraphs are increasing. It has been shown in [14] that we may assume further that $\Sigma_{\rho,h}$ is stable in the sense of MOTS.

Standard barriers for $\Sigma_{\rho,h}$ can be constructed from the trapped boundary ∂C_ρ by slight ($\mathcal{C}^2$)-inward perturbation above respectively below its boundary $\Gamma_{\rho,h}$, locally uniformly in (ρ, h), so that the angle (in the underlying Euclidean coordinate system) at which $\Sigma_{\rho,h}$ meets ∂C_ρ is bounded away from 0. For the special case of MOTS this inwards bending is explained in some detail in [12, Section 3]. Together with the λ-minimizing property and Allard's boundary regularity theorem [1] it follows that near ∂C_ρ the hypersurface $\Sigma_{\rho,h}$ can be written as a vertical $\mathcal{C}^{1,\alpha}$ graph above $\{x^n = h\}$. In particular, we see that $\Sigma_{\rho,h}$ intersects C_ρ transversely. Higher regularity of the defining function—which solves a prescribed mean curvature equation—then follows from Schauder theory in a standard way.

Finally, since each horizontal plane above $x^n = \Lambda$ has $\theta^+ > 0$ and each horizontal plane below $x^n = -\Lambda$ has $\theta^+ < 0$ by Lemma 5 it follows from Lemma 4 that each $\Sigma_{\rho,h}$ actually lies in $C_{\rho,\Lambda}$. □

Remark. We are grateful to Brian White for helping us clarify the following issue: When M is an n-dimensional $\mathcal{C}^{1,\alpha}$ manifold and g is a complete $\mathcal{C}^\alpha$ Riemannian metric on it, then the standard interior Allard-type $\mathcal{C}^{1,\alpha}$ regularity of (almost) minimizing boundaries away from a set of Hausdorff dimension at most 8 holds. This was shown by J. Taylor in [30] (this part of the discussion in her paper applies to n–dimensional manifolds). When the manifold is $\mathcal{C}^2$ and the metric is Lipschitz, then this follows also from the work of R. Schoen and L. Simon [24] (for almost minimizers this was pointed out by B. White in [32, p. 498]). When the manifold is $\mathcal{C}^4$ and the metric $\mathcal{C}^3$ so that the Nash embedding theorem provides an isometric embedding of (M, g) into a high dimensional Euclidean space, then this also follows directly upon applying the Euclidean regularity theory as in [29]. In the preceding lemma, note that once we know that our surfaces are $\mathcal{C}^{1,\alpha}$, we can then apply Schauder theory to the MOTS equation to obtain $\mathcal{C}^{3,\alpha}$ regularity. The boundary regularity follows more easily because the metric is conformal to the Euclidean metric there.

3.3. **Convergence of MOTS.** Although there is no reason to expect the family $\{\Sigma_{\rho,h}\}_{|h|\leq\Lambda}$ to be a $\mathcal{C}^{3,\alpha}$ foliation (even when there are no topological obstructions), we can still prove a partial regularity result. Its proof is similar to the proof of regularity of the outermost MOTS (established in [3] for $n = 3$ and then in [13] for $3 \leq n < 7$), but simpler because the two-sided λ-minimizing property eliminates concerns about embeddedness.

Lemma 7. *Let $\{\Sigma_{\rho,h}\}_{|h|\leq\Lambda}$ be as in Lemma 6. For each $h_0 \in (-\Lambda, \Lambda]$, the upper envelope of $\{\Sigma_{\rho,h}\}_{h<h_0}$ is a $\mathcal{C}^{3,\alpha}$ MOTS with boundary Γ_{ρ,h_0} which we denote by $\underline{\Sigma}_{\rho,h_0}$. By convention we define $\underline{\Sigma}_{\rho,-\Lambda} := \Sigma_{\rho,-\Lambda}$. Moreover, $\lim_{h\nearrow h_0} \Sigma_{\rho,h} = \underline{\Sigma}_{\rho,h}$ in the $\mathcal{C}^{3,\alpha}$ topology. We define $\overline{\Sigma}_{\rho,h_0}$ as the lower envelope of $\{\Sigma_{\rho,h}\}_{h>h_0}$ for $h_0 \in [-\Lambda, \Lambda)$ and $\overline{\Sigma}_{\rho,\Lambda} := \Sigma_{\rho,\Lambda}$ and note that analogous statements hold for these hypersurfaces.*

Proof. Fix $h_0 \in (\Lambda, \Lambda]$. Let $-\Lambda \leq h_i \nearrow h_0$ as $i \to \infty$ and pass the λ-minimizing boundaries Σ_{ρ,h_i} to a subsequential limit $\underline{\Sigma}_{\rho,h_0}$. Current convergence is automatic from the mass bounds, varifold convergence follows because there is no mass loss in limits of λ-minimizing currents, and $\mathcal{C}^{3,\alpha}$ convergence follows from Allard's interior and boundary regularity theorems. Note that since the Σ_{ρ,h_i} are increasing, this limit is independent of the choice of subsequence, and hence it is really a limit of the original sequence h_i. Moreover, the limit does not depend on choice of sequence h_i for the same reason. ☐

Definition 7. We say that $h_0 \in [-\Lambda, \Lambda]$ is a *jump height* if $\overline{\Sigma}_{\rho,h_0}$ does not equal $\underline{\Sigma}_{\rho,h_0}$.

By the previous lemma, $\Sigma_{\rho,h}$ converges to Σ_{ρ,h_0} in $\mathcal{C}^{3,\alpha}$ as $h \to h_0$ precisely when h_0 is *not* a jump height. It follows from the preceding lemma and Lemma 4 that h_0 is *not* a jump height if and only if the map $h \to \mathcal{L}^n(\Omega_{\rho,h})$ is continuous at h_0, where $\mathcal{L}^n(\Omega_{\rho,h})$ is the volume of the enclosed region $\Omega_{\rho,h}$ defined in the proof of Lemma 6. Note that this implies that there are at most countably many jump times $h_0 \in [-\Lambda, \Lambda]$.

We also observe that the MOTS constructed in Lemma 6 may be used to construct complete MOTS with good asymptotics.

Lemma 8. *For any choice of* $\rho_j \to \infty$ *and* $h_j \in [-\Lambda, \Lambda]$*, there exists a subsequence of* Σ_{ρ_j, h_j} *that converges in* $\mathcal{C}^{3,\alpha}$ *on compact subsets of* M *to a complete* $\mathcal{C}^{3,\alpha}$ *properly embedded MOTS* Σ_∞*. Moreover, there exists a constant* $c \in [-\Lambda, \Lambda]$ *such that outside a large compact subset of* M*,* Σ_∞ *can be written as the Euclidean graph* $\{x^n = f(x')\}$ *of some* $\mathcal{C}^{3,\alpha}$ *function* $f(x') = c + O^{3,\alpha}(|x'|^{3-n})$ *in the* $(x^1, \ldots, x^{n-1}, x^n) = (x', x^n)$ *coordinate system.*

Proof. Existence of a subsequential limit Σ_∞ and $\mathcal{C}^{3,\alpha}$ convergence follow as in the proof of Lemma 7. Note that the limit Σ_∞ is again λ-minimizing. Since each Σ_{ρ, h_ρ} lies between the horizontal planes $x^n = \pm\Lambda$, so does Σ_∞. The estimates below take part in the complement of a large ball B in M where we have harmonic asymptotics for the metric so that $g_{ij} = u^{\frac{4}{n-2}}\delta_{ij}$ where $u = 1 + O^{2,\alpha}(|x|^{2-n})$. It is not difficult to see from the corresponding property of Σ_{ρ_j} that the vertical projection of Σ_∞ onto the plane $\{x_n = 0\} \cap (M \setminus B)$ is surjective. Note that (7) implies that the Euclidean mean curvature of Σ_∞ is $O(|x|^{1-n})$. The λ-minimizing property of Σ_∞ gives rise to an explicit estimate of the form $O(|x'|^{-1})$ for the Euclidean area excess of Σ_∞ in large Euclidean balls centered at points $(x', x^n) \in \Sigma_\infty$ and of radius $|x'|/2$. Together with the Allard regularity theorem (the version in [29, Theorem 24.2] is particularly convenient here), this estimate implies that outside some large compact set, Σ_∞ is the graph of a function $f(x')$ such that $|f(x')| \leq \Lambda$ and $f(x') = O^{1,\gamma}(1)$ for some $\gamma \in (0, 1)$.

Since Σ_∞ is a MOTS we have that $H_{\Sigma_\infty} = -\operatorname{tr}_{\Sigma_\infty}(k)$. As in [22, p. 32], this translates into a prescribed (Euclidean) mean curvature equation for f via a conformal change (7). The initial estimate $f = O^{1,\gamma}(1)$, together with a computation of $\operatorname{tr}_{\Sigma_\infty} k$ as in the proof of Lemma 5, shows that the Euclidean mean curvature of f is $O^{\gamma}(|x'|^{1-n-\gamma})$ for some $\gamma \in (0, 1)$. Standard asymptotic analysis as in [20, 21] shows that there exists a constant $c \in [-\Lambda, \Lambda]$ such that $f(x') = c + O^{2,\gamma}(|x'|^{3-n})$. (Note that there is no logarithm term when $n - 1 = 2$ because f is bounded.) Repeating the above analysis with this information shows that $f(x') = c + O^{3,\alpha}(|x'|^{3-n})$, as asserted. □

Corollary 9. *When* $n > 3$*, the hypersurface* $\Sigma^{n-1}_\infty \subset M$ *in Lemma 8 is asymptotically flat and has zero energy with respect to its induced metric* g_∞*.*

The following lemma is a simple consequence of Proposition 3 and the fact that Σ_∞ is a limit of stable MOTS.

Lemma 10. *Let* Σ_∞ *be a complete MOTS whose existence is established by Lemma 8. For any* $v \in W^{1,2}_{\frac{3-n}{2}}(\Sigma_\infty)$*, we have*

$$\int_{\Sigma_\infty} \left(|\nabla v|^2 + Q_{\Sigma_\infty} v^2\right) \, d\mathcal{H}^{n-1} \geq 0.$$

We omit the proof because it is strictly simpler than that of Lemma 17 in Section 5.3. Specifically, the proof of Lemma 17 becomes a proof of Lemma 10 by simply replacing Z by an arbitrary compactly supported vector field on M and replacing the use of Lemma 15 by Proposition 3.

4. THE CASE $n = 3$

We consider the base case $n = 3$ of our inductive proof. After the preparation of the previous section, the rest of the proof of the $n = 3$ case is essentially the same as for the time-symmetric case in [25], where minimal surfaces are replaced by MOTS.

By Lemma 8, we can extract a subsequential limit Σ_∞ of $\Sigma_{\rho,0}$ as $\rho \to \infty$. Let Σ'_∞ be the noncompact component of Σ_∞. Lemma 10 implies that

$$\int_{\Sigma'_\infty} \left(|\nabla v|^2 + Q_{\Sigma'_\infty} v^2 \right) d\mathcal{H}^2 \geq 0 \tag{11}$$

for every $v \in W^{1,2}_{\frac{3-n}{2}}(\Sigma_\infty)$. Noting that Σ_∞ has quadratic area growth, we can use the "logarithmic cut-off trick" exactly as in [25, page 54] to approximate the constant function 1 on Σ'_∞ by compactly supported functions in order to conclude that

$$\int_{\Sigma'_\infty} Q_{\Sigma'_\infty} \, d\mathcal{H}^2 \geq 0.$$

The strict dominant energy condition then implies that

$$\int_{\Sigma'_\infty} K_{\Sigma'_\infty} \, d\mathcal{H}^2 > 0, \tag{12}$$

where $K_{\Sigma'_\infty}$ denotes the Gauss curvature. On the other hand, just as in [22], the estimate $(g_\infty)_{ij}(x') - \delta_{ij} = O^2(|x'|^{-1})$ implies that the geodesic curvature of $\partial(\Sigma'_\infty \cap C_r)$ is $\kappa = \frac{1}{r} + O(r^{-2})$ while the length of $\partial(\Sigma'_\infty \cap C_r)$ is $2\pi r + O(1)$. The Gauss-Bonnet Theorem tells us that

$$\int_{\Sigma'_\infty \cap C_r} K_{\Sigma'_\infty} \, d\mathcal{H}^2 = 2\pi\chi(\Sigma'_\infty \cap C_r) - \int_{\partial(\Sigma'_\infty \cap C_r)} \kappa \, d\mathcal{H}^1.$$

Combined with (12) and the asymptotics of $\partial(\Sigma'_\infty \cap C_r)$ described above, for large r, we obtain

$$0 < 2\pi\chi(\Sigma'_\infty \cap C_r) - 2\pi.$$

Since $\Sigma'_\infty \setminus C_r$ is a graph for large r, we know that $\Sigma'_\infty \cap C_r$ is connected, yielding a contradiction. $\square$

5. THE CASE $3 < n < 8$

Let $3 < n < 8$. We assume, in addition to the indirect assumption in Section 3 that Theorem 1 holds in $n-1$ dimensions. For the reasons described in the introduction, the proof here is substantially different from the proof in the time-symmetric case. The fundamental problem here is how to choose the MOTS $\Sigma_{\rho,h}$ with the appropriate height h. As seen in Section 4, this is not an issue when $n = 3$.

5.1. **The functional $\mathcal{F}$.**

In this section we consider a functional $\mathcal{F}$ that will be essential for our proof. In order to motivate the definition of $\mathcal{F}$, consider the time-symmetric case when $k = 0$ so that the MOTS $\{\Sigma_{\rho,h}\}_{|h|\leq\Lambda}$ constructed in Section 3.2 are minimal hypersurfaces. An important step in the proof of the Riemannian positive mass theorem when $3 < n < 8$ [22] is to pick Σ_{ρ,h_ρ} with least area in this family. Suppose for a moment that the family $\{\Sigma_{\rho,h}\}_{|h|\leq|\Lambda|}$ is actually a $\mathcal{C}^{3,\alpha}$ foliation of minimal hypersurfaces with a first-order deformation vector field X that is equal to ∂_n at ∂C_ρ. Then

$$\begin{aligned}
\frac{d}{dh}\mathcal{H}^{n-1}(\Sigma_{\rho,h}) &= \int_{\Sigma_{\rho,h}} (\operatorname{div}_{\Sigma_{\rho,h}} X)\, d\mathcal{H}^{n-1} \\
&= \int_{\Sigma_{\rho,h}} \operatorname{div}_{\Sigma_{\rho,h}}(\varphi\nu + \hat{X})\, d\mathcal{H}^{n-1} \\
&= \int_{\Sigma_{\rho,h}} (H\varphi + \operatorname{div}_{\Sigma_{\rho,h}} \hat{X})\, d\mathcal{H}^{n-1} \\
&= \int_{\Sigma_{\rho,h}} (\operatorname{div}_{\Sigma_{\rho,h}} \hat{X})\, d\mathcal{H}^{n-1} \\
&= \int_{\partial\Sigma_{\rho,h}} \langle \hat{X}, \eta\rangle\, d\mathcal{H}^{n-2} \\
&= \int_{\partial\Sigma_{\rho,h}} \langle \partial_n, \eta\rangle\, d\mathcal{H}^{n-2}.
\end{aligned}$$

If $h_\rho \in (-\Lambda, \Lambda)$ minimizes areas as described above, then the *first derivative* in h of the integral above is nonnegative. This, along with the stability of Σ_{ρ,h_ρ} among deformations that keep the boundary fixed, is all that is needed to finish the proof in the time-symmetric case [22].

We now return to the general case. Instead of using the area functional, which is not adapted for application to MOTS, we will build our proof around the functional described below.

Definition 8. Let Σ be a compact hypersurface in M whose boundary lies on some coordinate cylinder ∂C_r. Then we define

$$\mathcal{F}(\Sigma) = \int_{\partial\Sigma} \langle \partial_n, \eta\rangle \, d\mathcal{H}^{n-2}, \tag{13}$$

where η is the outward unit normal of $\partial\Sigma$ in Σ. Note that, using harmonic asymptotics, one can easily see that

$$\mathcal{F}(\Sigma) = \int_{\partial\Sigma} u^{\frac{2(n-1)}{n-2}} \eta_0^n \, d\mathcal{H}_0^{n-2}, \tag{14}$$

where η_0^n is the n-th component of the unit normal η_0 computed using the Euclidean metric, and $\mathcal{H}_0^{n-1}$ denotes Euclidean Hausdorff measure.

The barrier planes $\{x_n = \pm\Lambda\}$ give us a sign on $\mathcal{F}(\Sigma_{\rho,\pm\Lambda})$:

Lemma 11. *For any ρ sufficiently large,*

$$\mathcal{F}(\Sigma_{\rho,-\Lambda}) < 0 < \mathcal{F}(\Sigma_{\rho,\Lambda}).$$

Proof. From Lemma 6 we know that $\Sigma_{\rho,\Lambda}$ lies below the plane $\{x^n = \Lambda\}$ in C_ρ. The strong maximum principle Lemma 4 implies that they cannot meet tangentially at their common boundary $\Gamma_{\rho,\Lambda}$. Hence the Euclidean outward unit normal of $\partial\Sigma_{\rho,\Lambda}$ in $\Sigma_{\rho,\Lambda}$ satisfies $\eta_0^n > 0$. The inequality $\mathcal{F}(\Sigma_{\rho,\Lambda}) > 0$ then follows from (14). The proof that $\mathcal{F}(\Sigma_{\rho,-\Lambda}) < 0$ is analogous. ☐

Recall the definition of jump heights from Section 3.3.

Lemma 12. *The function $h \mapsto \mathcal{F}(\Sigma_{\rho,h})$ is continuous at every $h_0 \in [-\Lambda, \Lambda]$ that is* not *a jump height. If $h_0 \in [-\Lambda, \Lambda]$* is *a jump height, then*

$$\lim_{h\nearrow h_0} \mathcal{F}(\Sigma_{\rho,h}) \geq \mathcal{F}(\Sigma_{\rho,h_0}) \geq \lim_{h\searrow h_0} \mathcal{F}(\Sigma_{\rho,h}),$$

where both limits exist, and at least one of the inequalities above is strict. In other words, there must be a downward jump discontinuity at every jump height.

Proof. Let $h_0 \in [-\Lambda, \Lambda]$. Then by Lemma 7, $\lim_{h\nearrow h_0} \mathcal{F}(\Sigma_{\rho,h}) = \mathcal{F}(\underline{\Sigma}_{\rho,h_0})$ and $\lim_{h\searrow h_0} \mathcal{F}(\Sigma_{\rho,h}) = \mathcal{F}(\overline{\Sigma}_{\rho,h_0})$. By definition, if h_0 is not a jump height, then both of these limits must equal $\mathcal{F}(\Sigma_{\rho,h_0})$

Let h_0 be a jump height. Since the family $\{\Sigma_{\rho,h}\}_{|h|\leq\Lambda}$ is ordered, it is clear that $\underline{\Sigma}_{\rho,h_0}$ lies beneath Σ_{ρ,h_0}, which lies beneath $\overline{\Sigma}_{\rho,h_0}$. Since they all share the common boundary Γ_{ρ,h_0}, we have $(\underline{\eta})_0^n \geq \eta_0^n \geq (\overline{\eta})_0^n$, where $\underline{\eta}, \eta, \overline{\eta}$ are the outward normals of Γ_{ρ,h_0} in $\underline{\Sigma}_{\rho,h_0}$, Σ_{ρ,h_0}, and $\overline{\Sigma}_{\rho,h_0}$, respectively. Since h_0 is jump height, $\underline{\Sigma}_{\rho,h_0} \neq \overline{\Sigma}_{\rho,h_0}$, so the strong maximum principle (Lemma 4) implies that at least one of the above inequalities is strict. By the definition of $\mathcal{F}$ in (14),

$$\mathcal{F}(\underline{\Sigma}_{\rho,h_0}) \geq \mathcal{F}(\Sigma_{\rho,h_0}) \geq \mathcal{F}(\overline{\Sigma}_{\rho,h_0}),$$

where at least one of the inequalities is strict. ☐

Notation. Let Z be any $\mathcal{C}^2$ vector field on M that is equal to ∂_n outside some compact set. Given a hypersurface Σ equipped with a global unit normal ν, we use the decomposition

$$Z = \phi\nu + \hat{Z},$$

into normal and tangential components along Σ.

We now compute the first variation of $\mathcal{F}$. In view of Lemmas 11 and 12 we may hope to find $h_\rho \in (-\Lambda, \Lambda)$ such that the derivative of $h \to \mathcal{F}(\Sigma_{\rho,h})$ at h_ρ (defined in a suitably weak sense) is nonnegative.

Proposition 13. *Let Σ be a compact hypersurface in M whose boundary lies on some ∂C_r, equipped with a global unit normal ν. Let X be a $\mathcal{C}^1$ vector field along Σ that is tangent to ∂C_r at $\partial\Sigma$. As long as $Z = \partial_n$ along ∂C_r, we have*

$$D\mathcal{F}|_\Sigma(X) = \int_{\partial\Sigma} \langle \phi\nabla\varphi + G(X), \eta\rangle \, d\mathcal{H}^{n-2}, \tag{15}$$

where

$$G(X) = D_X Z - D_{\hat{Z}}\hat{X} + (\varphi H + \operatorname{div}_\Sigma \hat{X})\hat{Z} - \phi S(\hat{X}) - \varphi S(\hat{Z}). \tag{16}$$

Remark. Of course, it does not matter what Z is away from ∂C_r, and it is unnecessary to introduce Z at all except for the purposes of readability and for later application.

Proof. Note that $\mathcal{F}(\Sigma) = \int_{\partial\Sigma}\langle Z, \eta\rangle \, d\mathcal{H}^{n-2}$. Let $e_1, \ldots, e_{n-2}$ be a local orthonormal frame for $T(\partial\Sigma)$. We can differentiate Z, η, and the induced measure on $\partial\Sigma$ to obtain

$$D\mathcal{F}|_\Sigma(X) =$$

$$\int_{\partial\Sigma} \left[\langle D_X Z, \eta\rangle + \left\langle Z, \langle D_\eta X, \nu\rangle\nu - \sum_{i=1}^{n-2} \langle D_{e_i} X, \eta\rangle e_i \right\rangle + \langle Z, \eta\rangle \operatorname{div}_{\partial\Sigma} X \right] d\mathcal{H}^{n-2}. \tag{17}$$

(The derivative of η is computed by differentiating the orthogonality relations.) Along $\partial\Sigma$, we introduce the decomposition $\hat{Z} = \psi\eta + Z^\partial$ into components that are normal and tangential to $\partial\Sigma$. The second term in the integrand of (17) is

$$\begin{aligned} \left\langle Z, \langle D_\eta X, \nu\rangle\nu - \sum_{i=1}^{n-2} \langle D_{e_i} X, \eta\rangle e_i \right\rangle &= \langle Z, \nu\rangle\langle D_\eta X, \nu\rangle - \sum_{i=1}^{n-2} \langle Z, e_i\rangle\langle D_{e_i} X, \eta\rangle \\ &= \phi\langle D_\eta(\varphi\nu + \hat{X}), \nu\rangle - \langle D_{Z^\partial} X, \eta\rangle \\ &= \phi(\nabla_\eta\varphi + \langle D_\eta \hat{X}, \nu\rangle) - \langle D_{Z^\partial} X, \eta\rangle \\ &= \langle \phi\nabla\varphi, \eta\rangle - \langle \phi S(\hat{X}), \eta\rangle - \langle D_{Z^\partial} X, \eta\rangle. \end{aligned} \tag{18}$$

The third term in the integrand of (17) is

$$\begin{aligned}\langle Z,\eta\rangle \operatorname{div}_{\partial\Sigma} X &= \langle Z,\eta\rangle(\operatorname{div}_\Sigma X - \langle D_\eta X,\eta\rangle)\\ &= \langle \hat{Z},\eta\rangle(\varphi H + \operatorname{div}_\Sigma \hat{X}) - \psi\langle D_\eta X,\eta\rangle\\ &= \langle(\varphi H + \operatorname{div}_\Sigma \hat{X})\hat{Z},\eta\rangle - \langle D_{\psi\eta}X,\eta\rangle. \end{aligned} \tag{19}$$

Notice that the first term in the integrand of (17), the first two terms of (18) and the first term of (19) combine to give

$$\langle \phi\nabla\varphi + D_X Z + (\varphi H + \operatorname{div}_\Sigma \hat{X})\hat{Z} - \phi S(\hat{X}),\eta\rangle. \tag{20}$$

The remaining two terms, which are the last term of (18) and the last term of (19), combine to give

$$\begin{aligned}-\langle D_{Z^\partial}X,\eta\rangle - \langle D_{\psi\eta}X,\eta\rangle &= -\langle D_{\hat{Z}}X,\eta\rangle\\ &= -\langle D_{\hat{Z}}(\varphi\nu + \hat{X}),\eta\rangle\\ &= -\langle \varphi D_{\hat{Z}}\nu + D_{\hat{Z}}\hat{X},\eta\rangle\\ &= -\langle \varphi S(\hat{Z}) + D_{\hat{Z}}\hat{X},\eta\rangle. \end{aligned} \tag{21}$$

The result follows from combining (20) and (21). □

5.2. **Height picking and stability.** The following lemma, whose proof we defer to Section 5.4, will stand in for the geometric inequality (1) that was available in the time-symmetric case.

Lemma 14. *Let $\Sigma'_{\rho,h}$ denote the component of $\Sigma_{\rho,h}$ that contains the boundary $\Gamma_{\rho,h}$. For any large ρ, there exists $h_\rho \in (-\Lambda,\Lambda)$ and a $\mathcal{C}^2$ vector field X along Σ'_{ρ,h_ρ} that is equal to $Z = \partial_n$ at $\partial\Sigma'_{\rho,h_\rho} = \Gamma_{\rho,h_\rho}$, such that $\varphi = \langle X,\nu\rangle > 0$,*

$$D\theta^+|_{\Sigma'_{\rho,h_\rho}}(X) = 0, \tag{22}$$

and

$$D\mathcal{F}|_{\Sigma'_{\rho,h_\rho}}(X) \geq 0. \tag{23}$$

Consider $h \mapsto \Sigma_{\rho,h}$ as a "path" of $\mathcal{C}^{3,\alpha}$ hypersurfaces. *If* this path were differentiable in h (and if the $\Sigma_{\rho,h}$ were connected), then the proposition would be easy to prove. By Lemma 11, we could find h_ρ such that

$$\left.\frac{d}{dh}\mathcal{F}(\Sigma_{\rho,h})\right|_{h=h_\rho} \geq 0$$

and choose X to be the first-order deformation of the family $\Sigma_{\rho,h}$ at $h = h_\rho$. Then the inequality above would become (23), and the fact that each $\Sigma_{\rho,h}$ is a MOTS would imply that $D\theta^+|_{\Sigma_{\rho,h_\rho}}(X) = 0$. Unfortunately, $\Sigma_{\rho,h}$ need not be differentiable in h. In fact, if M has appropriate topology, the family $\Sigma_{\rho,h}$ must contain jumps. From Lemma 12 we know that $\mathcal{F}(\Sigma_{\rho,h})$ can only jump down at a jump height, so the presence of jumps does not cause any problem for finding h_ρ as described above. Even in the absence of jumps, the lack of differentiability in h presents a technical challenge.

Notation. For the remainder of this section, we will abbreviate Σ'_{ρ,h_ρ} by Σ_ρ.

Lemma 14 allows us to conclude the following stability-like property, which the reader should compare to Proposition 3.

Lemma 15. *Let ρ be sufficiently large, and choose X and φ as in the statement of Lemma 14. Then for any any $\mathcal{C}^1$ function v on Σ_ρ that is equal to $\phi = \langle Z, \nu\rangle$ at $\partial\Sigma_\rho$, we have*

$$\int_{\Sigma_\rho} (|\nabla v|^2 + Qv^2)\, d\mathcal{H}^{n-1} + \int_{\partial\Sigma_\rho} \langle \bar{G}(X), \eta\rangle\, d\mathcal{H}^{n-2} \geq 0 \tag{24}$$

where

$$\bar{G}(X) = G(X) + \phi\varphi W. \tag{25}$$

Proof. We begin by following the argument in [17]. Using equation (3), along with the positivity of φ, we have

$$\begin{aligned} D\theta^+|_{\Sigma_\rho}(X) &= -\Delta\varphi + 2\langle W, \nabla\varphi\rangle + (\operatorname{div} W - |W|^2 + Q)\varphi \\ &= -\Delta\varphi + (|\nabla\varphi|^2 + |W|^2 - |W - \nabla\varphi|^2)\varphi + (\operatorname{div} W - |W|^2 + Q)\varphi \\ &= -\Delta\varphi + |\nabla \log\varphi|^2\varphi - |W - \nabla\log\varphi|^2\varphi + (\operatorname{div} W + Q)\varphi \\ &= -(\Delta\log\varphi)\varphi - |W - \nabla\log\varphi|^2\varphi + (\operatorname{div} W + Q)\varphi \\ &= [\operatorname{div}(W - \nabla\log\varphi)]\varphi - |W - \nabla\log\varphi|^2\varphi + Q\varphi. \end{aligned}$$

Now let v be any $\mathcal{C}^1$ function on Σ_ρ that is equal to ϕ at $\partial\Sigma_\rho$. We multiply the above equation by $v^2\varphi^{-1}$ to obtain

$$\begin{aligned} v^2\varphi^{-1} D\theta^+|_{\Sigma_\rho}(X) &= [\operatorname{div}(W - \nabla\log\varphi)]v^2 - |W - \nabla\log\varphi|^2 v^2 + Qv^2 \\ &= \operatorname{div}(v^2(W - \nabla\log\varphi)) - \langle W - \nabla\log\varphi, 2v\nabla v\rangle \\ &\quad - |W - \nabla\log\varphi|^2 v^2 + Qv^2 \\ &= \operatorname{div}(v^2(W - \nabla\log\varphi)) \\ &\quad + |(W - \nabla\log\varphi)v|^2 + |\nabla v|^2 - |(W - \nabla\log\varphi)v + \nabla v|^2 \\ &\quad - |W - \nabla\log\varphi|^2 v^2 + Qv^2 \\ &= \operatorname{div}(v^2(W - \nabla\log\varphi)) + |\nabla v|^2 + Qv^2 \\ &\quad - |(W - \nabla\log\varphi)v + \nabla v|^2. \end{aligned} \tag{26}$$

Together with equation (22), this implies

$$0 \leq |\nabla v|^2 + Qv^2 + \operatorname{div}(v^2(W - \nabla \log \varphi)).$$

Noting that $v = \phi = \varphi$ at $\partial\Sigma_\rho$, we can integrate this to obtain

$$\begin{aligned}
0 &\leq \int_{\Sigma_\rho} (|\nabla v|^2 + Qv^2)\, d\mathcal{H}^{n-1} + \int_{\partial\Sigma_\rho} \langle v^2(W - \nabla \log \varphi), \eta\rangle \, d\mathcal{H}^{n-2} \\
&= \int_{\Sigma_\rho} (|\nabla v|^2 + Qv^2)\, d\mathcal{H}^{n-1} + \int_{\partial\Sigma_\rho} \langle \phi\varphi W - \phi\nabla\varphi, \eta\rangle \, d\mathcal{H}^{n-2} \\
&= \int_{\Sigma_\rho} (|\nabla v|^2 + Qv^2)\, d\mathcal{H}^{n-1} + \int_{\partial\Sigma_\rho} \langle \bar{G}(X), \eta\rangle \, d\mathcal{H}^{n-2} - D\mathcal{F}|_{\Sigma_\rho}(X) \\
&\leq \int_{\Sigma_\rho} (|\nabla v|^2 + Qv^2)\, d\mathcal{H}^{n-1} + \int_{\partial\Sigma_\rho} \langle \bar{G}(X), \eta\rangle \, d\mathcal{H}^{n-2},
\end{aligned}$$

where we used (15) and (23) in the third and fourth lines, respectively. □

5.3. **Analysis of the complete MOTS limit.** The strategy for the rest of the proof of Theorem 1 follows that of the time-symmetric case treated in [22] closely. By Lemma 8, there exists a sequence $\rho_j \to \infty$ such that Σ_{ρ_j} converges locally in $\mathcal{C}^{3,\alpha}$ to some Σ_∞ that has the properties described in Lemma 8.

While Lemma 10 is strong enough to show that Σ_∞ is conformal to a scalar-flat asymptotically flat manifold, it is not strong enough to control the change of mass under this conformal change. For that, we need to be able to allow test functions that are asymptotic to 1. In order to prove this, we will need to have uniform control over the Σ_ρ's as in the following lemma.

Lemma 16. *Recall the definitions of Z and ϕ from Section 5.1. The following estimate holds uniformly in ρ.*

$$\left(\int_{\Sigma_\rho\smallsetminus C_r} \left(|\nabla\phi|^2 + Q_{\Sigma_\rho}\phi^2\right)\, d\mathcal{H}^{n-1} + \int_{\partial(\Sigma_\rho\smallsetminus C_r)} \langle \bar{G}(Z), \eta\rangle \, d\mathcal{H}^{n-2}\right) = O(r^{-1}). \tag{27}$$

Proof. We would first like to show that the estimates

$$D\theta^+|_{\Sigma_\rho}(Z) = O(|x|^{-n}) \tag{28}$$

$$D\mathcal{F}|_{\Sigma_\rho\cap C_r}(Z) = O(r^{-1}) \tag{29}$$

hold uniformly in ρ. Using harmonic asymptotics, formula (7) applies outside of a compact set to give us

$$H = u^{\frac{-2}{n-2}}\left(H_0 + \tfrac{2(n-1)}{n-2} u^{-1}\nabla_{\nu_0} u\right),$$

where H_0 and ν_0 are the mean curvature and upward unit normal with respect to the Euclidean metric. Clearly, H_0 and ν_0 do not change under vertical translation, and H_0 is bounded, therefore the decay of u implies that $DH|_{\Sigma_\rho}(Z) = O(|x|^{-n})$. The $\operatorname{tr}_{\Sigma_\rho} k$ term is easier to handle, giving us (28). To derive (29), we simply use (14), the fact that ν_0 does not change under vertical translation, and decay of u.

To prove the lemma, we vary Σ_ρ in the direction Z, which is just vertical translation outside a compact set. We repeat the exact same argument from the first part of Lemma 15, except we use the region $\Sigma_\rho \smallsetminus C_r$ instead of Σ_ρ, the vector field Z instead of X, the function ϕ instead of φ, and we take $v = \phi$. Then on $\Sigma_\rho \smallsetminus C_r$, equation (26) becomes

$$\phi D\theta^+|_{\Sigma_\rho}(Z) = \operatorname{div}(\phi^2 W - \phi\nabla\phi) + |\nabla\phi|^2 + Q\phi^2 - |W|^2\phi^2.$$

Note that ϕ need not be positive in order for this to hold. Therefore

$$\begin{aligned}
&\int_{\Sigma_\rho\smallsetminus C_r} (|\nabla\phi|^2 + Q\phi^2)\, d\mathcal{H}^{n-1} + \int_{\partial(\Sigma_\rho\smallsetminus C_r)} \langle \bar{G}(Z), \eta\rangle\, d\mathcal{H}^{n-2} \\
&\quad = \int_{\Sigma_\rho\smallsetminus C_r} (|\nabla\phi|^2 + Q\phi^2)\, d\mathcal{H}^{n-1} + \int_{\partial(\Sigma_\rho\smallsetminus C_r)} \langle \phi^2 W - \phi\nabla\phi, \eta\rangle\, d\mathcal{H}^{n-2} \\
&\qquad + D\mathcal{F}|_{\Sigma_\rho}(Z) - D\mathcal{F}|_{\Sigma_\rho\cap C_r}(Z) \\
&\quad = \int_{\Sigma_\rho\smallsetminus C_r} \left(\phi D\theta^+|_{\Sigma_\rho}(Z) + |W|^2\phi^2\right)\, d\mathcal{H}^{n-2} \\
&\quad = O(r^{-1}),
\end{aligned}$$

where the last line follows from the (28) and (29), decay of W, and volume control coming from the λ-minimizing property of Σ_ρ. ☐

The following lemma is the analytical consequence of our careful height-picking in Section 5.2.

Lemma 17. *Let v be a function on Σ_∞ such that $v - 1 \in W^{1,2}_{\frac{3-n}{2}}(\Sigma_\infty)$. Then*

$$\int_{\Sigma_\infty} \left(|\nabla v|^2 + Q_{\Sigma_\infty} v^2\right)\, d\mathcal{H}^{n-1} \geq 0. \tag{30}$$

Proof. Following the notation of Lemma 8 we use coordinates x' on $\Sigma_\infty \smallsetminus B$ where B is a large compact subset of M. It is easy to see from Lemma 8 that

$$Q_{\Sigma_\infty} = \tfrac{1}{2}R_{\Sigma_\infty} - \mu - J(\nu_{\Sigma_\infty}) - \tfrac{1}{2}|k_{\Sigma_\infty} + B_{\Sigma_\infty}|^2 = O(|x'|^{-n}).$$

Since the volume growth of Σ_∞ is like that of $\mathbb{R}^{n-1}$, one can then see that

$$\int_{\Sigma_\infty} \left(|\nabla v|^2 + |Q_{\Sigma_\infty}| v^2\right)\, d\mathcal{H}^{n-1} < \infty, \tag{31}$$

whenever $v - 1 \in W^{1,2}_{\frac{3-n}{2}}(\Sigma_\infty)$.

We will write Σ_j for $\Sigma_{\rho_j} = \Sigma'_{\rho_j, h_{\rho_j}}$ in this proof and use a j-subscript to denote quantities that depend on Σ_j, including $j = \infty$. Whenever $r < \rho_j$, we use η_j to denote the outward unit normal of $\partial(\Sigma_j \cap C_r)$ inside Σ_j. (In what follows, we can restrict our attention to values of r such that ∂C_r is transverse to each Σ_j.) Fix a $\mathcal{C}^2$ vector field Z on M that agrees with ∂_n outside a compact set. Let $\phi_\infty = \langle \nu_{\Sigma_\infty}, Z\rangle$ and $\phi_j = \langle \nu_{\Sigma_j}, Z\rangle$. Note that Lemma 8 implies that $\phi_\infty - 1 = O^1(|x'|^{2-n}) \in W^{1,2}_{\frac{3-n}{2}}(\Sigma_\infty)$ and that $\bar{G}_\infty(Z) = O(|x'|^{1-n})$ where $\bar{G}_\infty$ is as (25). We have that

$$\begin{aligned}
&\int_{\Sigma_\infty} \left(|\nabla\phi_\infty|^2 + Q_\infty \phi_\infty^2\right)\, d\mathcal{H}^{n-1} \\
&= \lim_{r\to\infty} \int_{\Sigma_\infty \cap C_r} \left(|\nabla\phi_\infty|^2 + Q_\infty \phi_\infty^2\right)\, d\mathcal{H}^{n-1} \\
&= \lim_{r\to\infty} \left(\int_{\Sigma_\infty \cap C_r} \left(|\nabla\phi_\infty|^2 + Q_\infty \phi_\infty^2\right)\, d\mathcal{H}^{n-1} + \int_{\partial(\Sigma_\infty \cap C_r)} \langle \bar{G}_\infty(Z), \eta_\infty\rangle \, d\mathcal{H}^{n-2} \right) \\
&= \lim_{r\to\infty} \lim_{j\to\infty} \left(\int_{\Sigma_j \cap C_r} \left(|\nabla\phi_j|^2 + Q_j \phi_j^2\right)\, d\mathcal{H}^{n-1} + \int_{\partial(\Sigma_j \cap C_r)} \langle \bar{G}_j(Z), \eta_j\rangle \, d\mathcal{H}^{n-2} \right) \\
&= \lim_{j\to\infty} \left(\int_{\Sigma_j} \left(|\nabla\phi_j|^2 + Q_j \phi_j^2\right)\, d\mathcal{H}^{n-1} + \int_{\partial\Sigma_j} \langle \bar{G}_j(Z), \eta_j\rangle \, d\mathcal{H}^{n-2} \right) \\
&\geq 0,
\end{aligned}$$

where the last inequality follows from Lemma 15, and the fourth equality follows from Lemma 16. Hence (30) holds for $v = \phi_\infty$. Moreover, since the argument works for *any* Z that equals ∂_n outside a compact set, (30) holds for all test functions that agree with ϕ_∞ outside a compact set.

We now argue by density that the lemma holds for any function v such that $v - \phi_\infty \in W^{1,2}_{\frac{3-n}{2}}(\Sigma_\infty)$. Note that $\mathcal{C}^{3,\alpha}_c(\Sigma_\infty)$ is dense in $W^{1,2}_{\frac{3-n}{2}}(\Sigma_\infty)$. Let $v_i - \phi_\infty$ be a sequence of functions in $\mathcal{C}^{3,\alpha}_c(\Sigma_\infty)$ that converges to $v - \phi_\infty$ in $W^{1,2}_{\frac{3-n}{2}}(\Sigma_\infty)$. It is now straightforward to check that

$$\begin{aligned}
0 &\leq \liminf_{i\to\infty} \int_{\Sigma_\infty} \left(|\nabla v_i|^2 + Q_\infty v_i^2\right)\, d\mathcal{H}^{n-1} \\
&= \liminf_{i\to\infty} \int_{\Sigma_\infty} \big[(|\nabla v|^2 + Q_\infty v^2) + (2\nabla v\cdot\nabla(v_i - v) + 2Q_\infty v(v_i - v)) \\
&\quad +(|\nabla(v_i - v)|^2 + Q_\infty(v_i - v)^2)\big]\, d\mathcal{H}^{n-1} \\
&= \int_{\Sigma_\infty} \left(|\nabla v|^2 + Q_\infty v^2\right)\, d\mathcal{H}^{n-1},
\end{aligned}$$

where the cross terms vanish because of (31). Finally, we note that $\phi_\infty - 1 \in W^{1,2}_{\frac{3-n}{2}}$ to obtain the desired result. □

We are now ready to derive a contradiction to our induction hypothesis. Without loss of generality, we throw away all compact components of Σ_∞, noting that Lemmas 10 and 17 still hold on the remaining asymptotically flat component. The strict dominant energy condition implies that $Q_\infty < \frac{1}{2}R_{\Sigma_\infty}$, so Lemma 10 implies that for any nonzero $v \in W^{1,2}_{\frac{3-n}{2}}(\Sigma_\infty)$, we have

$$\int_{\Sigma_\infty} \left(|\nabla v|^2 + \tfrac{1}{2}R_{\Sigma_\infty} v^2\right)\, d\mathcal{H}^{n-1} > 0.$$

Using the fact that $n > 3$, we have

$$\int_{\Sigma_\infty} \left(|\nabla v|^2 + \tfrac{n-3}{4(n-2)}R_{\Sigma_\infty} v^2\right)\, d\mathcal{H}^{n-1} > 0, \tag{32}$$

for all nonzero $v \in W^{1,2}_{\frac{3-n}{2}}(\Sigma_\infty)$. This inequality implies that the conformal Laplacian

$$\Delta_{\Sigma_\infty} - \tfrac{n-3}{4(n-2)}R_{\Sigma_\infty} : W^{1,2}_{\frac{3-n}{2}}(\Sigma_\infty) \longrightarrow W^{-1,2}_{\frac{-1-n}{2}}(\Sigma_\infty)$$

is an isomorphism. In particular, we can find some nonzero $v \in W^{1,2}_{\frac{3-n}{2}}(\Sigma_\infty)$ such that

$$\Delta_{\Sigma_\infty} v - \tfrac{n-3}{4(n-2)}R_{\Sigma_\infty} v = \tfrac{n-3}{4(n-2)}R_{\Sigma_\infty}.$$

Setting $w = 1 + v$, we have

$$\Delta_{\Sigma_\infty} w - \tfrac{n-3}{4(n-2)}R_{\Sigma_\infty} w = 0.$$

By elliptic regularity, $w \in \mathcal{C}^{2,\alpha}_{loc}$, and by [20, Theorem 2], we must have $w(x') = 1 + O^{2,\alpha}(|x'|^{3-n})$. In particular, if g_∞ is the induced metric on Σ_∞, then by applying (6) in dimension $n-1$, the conformal metric $w^{\frac{4}{n-3}} g_\infty$ on Σ_∞^{n-1} is an asymptotically flat Riemannian manifold of zero scalar curvature. Using (8) in dimension $n-1$, the new zero scalar curvature metric on Σ_∞ has energy

$$\begin{aligned} E(w^{\frac{4}{n-3}}g_\infty) &= E(g_\infty) - \tfrac{2}{(n-3)\omega_{n-2}} \lim_{r\to\infty} \int_{\partial(\Sigma_\infty\cap C_r)} w\nabla_\nu w \, d\mathcal{H}^{n-1} \\ &= 0 - \tfrac{2}{(n-3)\omega_{n-2}} \int_{\Sigma_\infty} (|\nabla w|^2 + w\Delta w)\, d\mathcal{H}^{n-1} \\ &= \tfrac{-2}{(n-3)\omega_{n-2}} \int_{\Sigma_\infty} \left(|\nabla w|^2 + \tfrac{n-3}{4(n-2)} R_{\Sigma_\infty} w^2\right) d\mathcal{H}^{n-1}. \end{aligned}$$

Now since $w-1 \in W^{1,2}_{\frac{3-n}{2}}(\Sigma_\infty)$ and is not zero, we can use Lemma 17 plus the short argument used to derive (32) to see that

$$\int_{\Sigma_\infty} \left(|\nabla w|^2 + \tfrac{n-3}{4(n-2)} R_{\Sigma_\infty} w^2\right) d\mathcal{H}^{n-1} > 0,$$

and hence $E(w^{\frac{4}{n-3}}g_\infty) < 0$. This contradicts the time-symmetric case of Theorem 1 in dimension $n-1$, which is our induction hypothesis.

5.4. **Proof of Lemma 14.**

We define

$$h_\rho = \inf\{h \in [-\Lambda, \Lambda] \,|\, \mathcal{F}(\Sigma_{\rho,h}) > 0\}. \tag{33}$$

Our goal for this section is to prove that this choice of h_ρ satisfies the conclusion of Lemma 14. For now let us assume that Σ_{ρ,h_ρ} is connected.

Using Lemmas 11 and 12 it is easy to see that h_ρ exists, lies in $(-\Lambda,\Lambda)$, is not a jump height, and that $\mathcal{F}(\Sigma_{\rho,h_\rho}) = 0$. Although the path $h \to \Sigma_{\rho,h}$ of $\mathcal{C}^{3,\alpha}$ hypersurfaces need not be differentiable in h at h_ρ, the continuity of the map $h \mapsto \Sigma_{\rho,h}$ at $h = h_\rho$ allows us to use the inverse function theorem to *describe* the family $\Sigma_{\rho,h}$ precisely for h near h_ρ, as we will see below. We note that the work of B. White [31] on the moduli space of minimal submanifolds with boundary in $\mathbb{R}^n$ is useful for understanding the family $\Sigma_{\rho,h}$. However, since we do not need the full power of [31], we choose to use a simpler approach similar to that of [15] in our Case 2 below.

Since Σ_{ρ,h_ρ} is transverse to ∂C_ρ, we can find a $\mathcal{C}^{2,\alpha}$ vector field ∂_τ such that $\langle \partial_\tau, \nu\rangle > 0$ along Σ_{ρ,h_ρ}, and $\partial_\tau = Z = \partial_n$ at Γ_{ρ,h_ρ}. By integrating ∂_τ, there exists a relatively open neighborhood U of Σ_{ρ,h_ρ} in C_ρ and a $\mathcal{C}^{2,\alpha}$ diffeomorphism $F : \Sigma_{\rho,h_\rho} \times (-\delta,\delta) \longrightarrow U$ such that $F(\cdot,0)$ is the identity map, and $F_*(\frac{\partial}{\partial\tau}) = \partial_\tau$. In particular, $F(\cdot,\tau)$ maps $\partial\Sigma_{\rho,h_\rho} = \Gamma_{\rho,h_\rho}$ to $\Gamma(h_\rho+\tau)$ by vertical translation.

Any $\mathcal{C}^{2,\alpha}$ function $w : \Sigma_{\rho,h_\rho} \longrightarrow (-\delta,\delta)$ defines a graph in $\Sigma_{\rho,h_\rho} \times (-\delta,\delta)$, whose image under the diffeomorphism F is a $\mathcal{C}^{2,\alpha}$ hypersurface in U which we denote graph$[w]$, whose boundary lies on ∂C_ρ. Since h_ρ is not a jump height, Lemma 7 tells us that for each h sufficiently close to h_ρ, each $\Sigma_{\rho,h} = \mathrm{graph}[w]$ for some $\mathcal{C}^{2,\alpha}$ function w. Moreover, the function w converges to 0 in $\mathcal{C}^{2,\alpha}$ as h approaches h_ρ.

Recall that the operator $L_{\Sigma_{\rho,h_\rho}}$ on Σ_{ρ,h_ρ} defined by (4) has a principal Dirichlet eigenvalue, which is nonnegative since Σ_{ρ,h_ρ} is stable. We consider two cases, depending on whether this principal eigenvalue is positive or zero.

Case 1: The principal eigenvalue of $L_{\Sigma_{\rho,h_\rho}}$ is positive.

Define the map[3] $\Psi : \mathcal{C}^{2,\alpha}_0(\Sigma_{\rho,h_\rho}) \times \mathbb{R} \longrightarrow \mathcal{C}^{0,\alpha}(\Sigma_{\rho,h_\rho}) \times \mathbb{R}$ by

$$\Psi(w, s) = (\theta^+_{\mathrm{graph}[w+s]}, s).$$

Using (3) and (4), we see that

$$D\Psi|_{(0,0)}(w', s') = (L_{\Sigma_{\rho,h_\rho}}\langle (w' + s')\partial_\tau, \nu\rangle, s').$$

Since $L_{\Sigma_{\rho,h_\rho}} : \mathcal{C}^{2,\alpha}_0 \longrightarrow \mathcal{C}^{0,\alpha}$ is an isomorphism, so is $D\Psi|_{(0,0)}$. The inverse function theorem tells us that there exists a $\mathcal{C}^1$ function $\Phi : (-\epsilon, \epsilon) \longrightarrow \mathcal{C}^{2,\alpha}_0(\Sigma_{\rho,h_\rho})$ such that $\Phi(s)$ is the unique small solution of the equation $\Psi(\Phi(s), s) = (0, s)$. That is, it is the unique $\mathcal{C}^{2,\alpha}$-nearby MOTS whose boundary is $\Gamma_{\rho,h_\rho+s}$. So by continuity of $\Sigma_{\rho,h}$ at $h = h_\rho$ and the uniqueness, we must have that $\Sigma_{\rho,h_\rho+s} = \mathrm{graph}[\Phi(s) + s]$ for small s. Therefore $h \mapsto \Sigma_{\rho,h}$ is actually a $\mathcal{C}^1$ map into the space of $\mathcal{C}^{2,\alpha}$ hypersurfaces at $h = h_\rho$, so that the straightforward argument described in Section 5.2 applies. Specifically, if we take $X = (\frac{d\Phi}{ds}+1)\partial_\tau \in \mathcal{C}^{2,\alpha}$, then X is a first-order deformation vector field for the family $\Sigma_{\rho,h}$. Since each $\Sigma_{\rho,h}$ is a MOTS, $D\theta^+|_{\Sigma_{\rho,h_\rho}}(X) = \frac{d}{dh}\theta^+_{\Sigma_{\rho,h}}|_{h=h_\rho} = 0$. By construction of h_ρ, we have $D\mathcal{F}|_{\Sigma_{\rho,h_\rho}}(X) = \frac{d}{dh}\mathcal{F}(\Sigma_{\rho,h})\big|_{h=h_\rho} \geq 0$. Finally, the normal component φ of X is nonnegative on $\Sigma_{\rho,h}$ and positive on $\Gamma_{\rho,h}$ by construction, and hence positive everywhere by the strong maximum principle applied to the linear equation $L_{\Sigma_{\rho,h_\rho}}\varphi = 0$. This completes the proof of Lemma 14 for Case 1.

Case 2: The principal eigenvalue of $L_{\Sigma_{\rho,h_\rho}}$ is zero.

In this case, since Σ_{ρ,h_ρ} is connected, $L_{\Sigma_{\rho,h_\rho}}$ has a one-dimensional kernel in $\mathcal{C}^{2,\alpha}_0$ that is generated by a function that is positive away from the boundary. The same is then true for the adjoint $L^*_{\Sigma_{\rho,h_\rho}}$. We define the map $\Psi : \mathcal{C}^{2,\alpha}_0(\Sigma_{\rho,h_\rho})\times\mathbb{R}^2 \longrightarrow \mathcal{C}^{0,\alpha}(\Sigma_{\rho,h_\rho})\times\mathbb{R}^2$ by

$$\Psi(w, \kappa, s) = \left(-\kappa + \theta^+_{\mathrm{graph}[w+s]}, V(\mathrm{graph}[w + s]), s\right),$$

---

[3] Technically, it is only defined on some neighborhood of the origin.

where $V(\Sigma)$ denotes the "signed volume" of the region of C_ρ lying above Σ_{ρ,h_ρ} and below Σ. Then

$$D\Psi|_{(0,0,0)}(w',\kappa',s') = \left(L_{\Sigma_{\rho,h_\rho}}\langle (w'+s')\partial_\tau, \nu\rangle - \kappa', \int_{\Sigma_{\rho,h_\rho}} \langle (w'+s')\partial_\tau, \nu\rangle \, d\mathcal{H}^{n-1}, s' \right).$$

We claim that $D\Psi|_{(0,0,0)}$ is injective. If $D\Psi|_{(0,0,0)}(w',\kappa',s') = (0,0,0)$, then obviously $s' = 0$, and since the image of $\mathcal{C}^{2,\alpha}_0$ under $L_{\Sigma_{\rho,h_\rho}}$ is orthogonal to the kernel of $L^*_{\Sigma_{\rho,h_\rho}}$, $\kappa' = 0$. Finally, $\langle w'\partial_\tau, \nu\rangle$ is in the kernel of $L_{\Sigma_{\rho,h_\rho}}$ and has zero integral, and is therefore zero, proving the claim. Since $D\Psi|_{(0,0,0)}$ has index zero, it is also an isomorphism. So the inverse function theorem tells us that there exists a $\mathcal{C}^1$ function

$$(\Phi_1, \Phi_2) : (\epsilon, \epsilon) \times (\epsilon, \epsilon) \longrightarrow \mathcal{C}^{2,\alpha}_0(\Sigma_{\rho,h_\rho}) \times \mathbb{R},$$

such that $(\Phi_1(\xi,s), \Phi_2(\xi,s))$ is the unique small solution of the equation

$$\Psi(\Phi_1(\xi,s), \Phi_2(\xi,s), s) = (0,\xi,s).$$

In particular, graph$[\Phi_1(\xi,s)+s]$ is the unique $\mathcal{C}^{2,\alpha}$-nearby constant θ^+ hypersurface whose boundary is $\Gamma_{\rho,h_\rho+s}$ and whose "signed volume" is ξ. Define $\xi(s) := V(\Sigma_{\rho,h_\rho+s})$. By continuity of $\Sigma_{\rho,h}$ at $h = h_\rho$ and the uniqueness, it must be the case that $\Sigma_{\rho,h_\rho+s} =$ graph$[\Phi_1(\xi(s),s)+s]$ for small s. Note that since $\Sigma_{\rho,h_\rho+s}$ lies strictly above Σ_{ρ,h_ρ}, the function $\Phi_1(\xi(s),s)+s$ must be positive. The complication here is that we do not know that $\xi(s)$ depends on s in any nice way. Below, we will see that this does not matter.

By construction of h_ρ, we can choose a sequence $s_k \searrow 0$ such that

$$\mathcal{F}(\Sigma_{\rho,h_\rho+s_k}) > 0.$$

Let us pass to a subsequence such that the unit vector in the direction $(\xi(s_k), s_k)$ converges. We consider two subcases.

Case 2a: The ratio $\frac{\xi(s_k)}{s_k}$ converges to some finite number as $k \to \infty$.

The basic idea here is that even though we cannot take derivatives as in Case 1, we can take subsequential limits of difference quotients instead. The hypothesis of this subcase implies that $\frac{1}{s_k}[\Phi_1(\xi(s_k),s_k)+s_k]$ converges to some $\mathcal{C}^{2,\alpha}$ function $\bar\varphi$. Therefore we can take the limit of $\frac{1}{s_k}\theta^+_{\text{graph}[\Phi_1(\xi(s_k),s_k)+s_k]} = 0$ to obtain

$$L_{\Sigma_{\rho,h_\rho}}(\langle \bar\varphi\partial_\tau, \nu\rangle) = 0.$$

We claim that the conclusion of Lemma 14 holds for $X = \bar\varphi\partial_\tau \in \mathcal{C}^{2,\alpha}$. The previous equation tells us that $D\theta^+|_{\Sigma_{\rho,h_\rho}}(X) = 0$, and we can see that the normal component

$\varphi = \bar{\varphi}\langle\partial_\tau, \nu\rangle$ is nonnegative and equal to $\langle Z, \nu\rangle > 0$ at Γ_{ρ,h_ρ}. By the strong maximum principle for $L_{\Sigma_{\rho,h_\rho}}$, φ is strictly positive. Finally, since $\mathcal{F}(\Sigma_{\rho,h_\rho+s_k}) > 0$,

$$\begin{aligned} D\mathcal{F}|_{\Sigma_{\rho,h_\rho}}(X) &= \lim_{k\to\infty} \frac{1}{s_k}[\mathcal{F}(\text{graph}[\Phi_1(\xi(s_k), s_k) + s_k]) - \mathcal{F}(\Sigma_{\rho,h_\rho})] \\ &= \lim_{k\to\infty} \frac{1}{s_k}[\mathcal{F}(\Sigma_{\rho,h_\rho+s_k}) - \mathcal{F}(\Sigma_{\rho,h_\rho})] \\ &\geq 0. \end{aligned}$$

Case 2b: The ratio $\frac{s_k}{\xi(s_k)}$ converges to zero as $k \to \infty$.

We will show that this is impossible. The assumption implies that $\frac{1}{\xi(s_k)}[\Phi_1(\xi(s_k), s_k) + s_k]$ converges to some $\mathcal{C}^{2,\alpha}$ function $\bar{\varphi}$ that is zero at Γ_{ρ,h_ρ}. As in Case 2a, we conclude that

$$L_{\Sigma_{\rho,h_\rho}}(\langle\bar{\varphi}\partial_\tau, \nu\rangle) = 0.$$

Let $X = \bar{\varphi}\partial_\tau \in \mathcal{C}^{2,\alpha}$. Again, its normal component $\varphi = \bar{\varphi}\langle\partial_\tau, \nu\rangle$ is nonnegative, but this time it has zero boundary values. Since we also have $L_{\Sigma_{\rho,h_\rho}}\varphi = 0$, the strong maximum principle tells us that either φ is identically zero, or else $\nabla_\eta\varphi < 0$ at Γ_{ρ,h_ρ}. To see why φ cannot be zero, take the limit of the equation $\frac{V(\Sigma_{\rho,h_\rho+s_k})}{\xi(s_k)} = 1$ to obtain $\int_{\Sigma_{\rho,h_\rho}} \varphi \, d\mathcal{H}^{n-1} = 1$. Just as in Case 2a,

$$\begin{aligned} D\mathcal{F}|_{\Sigma_{\rho,h_\rho}}(X) &= \lim_{k\to\infty} \frac{1}{\xi(s_k)}[\mathcal{F}(\text{graph}[\Phi_1(\xi(s_k), s_k) + s_k]) - \mathcal{F}(\Sigma_{\rho,h_\rho})] \\ &= \lim_{k\to\infty} \frac{1}{\xi(s_k)}[\mathcal{F}(\Sigma_{\rho,h_\rho+s_k}) - \mathcal{F}(\Sigma_{\rho,h_\rho})] \\ &\geq 0. \end{aligned}$$

On the other hand, note that since $X = 0$ along Γ_{ρ,h_ρ}, only the second term in the identity (17) contributes to $D\mathcal{F}|_{\Sigma_{\rho,h_\rho}}$ so that

$$\begin{aligned} D\mathcal{F}|_{\Sigma_{\rho,h_\rho}}(X) &= \int_{\Gamma_{\rho,h_\rho}} \langle Z, \langle D_\eta X, \nu\rangle\nu\rangle d\mathcal{H}^{n-2} \\ &= \int_{\Gamma_{\rho,h_\rho}} \phi\nabla_\eta\varphi \, d\mathcal{H}^{n-2} \\ &< 0 \end{aligned}$$

since $\nabla_\eta\varphi < 0$ and $\phi > 0$ on Γ_{ρ,h_ρ}. This contradiction shows that Case 2b is impossible, finishing the proof of Lemma 14 when Σ_{ρ,h_ρ} is connected.

Now suppose Σ_{ρ,h_ρ} is not connected, and let $\Sigma'_{\rho,h}$ denote the component of $\Sigma_{\rho,h}$ that contains the boundary. Observe that since h_ρ is not a jump height, $\Sigma'_{\rho,h}$ converges to Σ'_{ρ,h_ρ} in $\mathcal{C}^{3,\alpha}$ as $h \to h_\rho$, and also that $\mathcal{F}(\Sigma'_{\rho,h}) = \mathcal{F}(\Sigma_{\rho,h})$ for all $h \in [-\Lambda, \Lambda]$. Using these two facts, we can see that the entire argument above can be executed with Σ'_{ρ,h_ρ} in place of $\Sigma_{\rho,h}$.

6. The density theorem

Recall that $\pi = k - (\operatorname{tr}_g k)g$. In this section it will be convenient to formulate everything in terms of π rather than k. With a slight abuse of terminology, we will refer to (M, g, π) as an initial data set. The purpose of this section is to prove the following theorem.

Theorem 18. *Let (M^n, g, π) be an asymptotically flat initial data set of type (p, q, q_0, α), as defined in Section 2. Assume that the dominant energy condition $\mu \geq |J|_g$ holds. For every $\epsilon > 0$ there exists asymptotically flat initial data $(\bar{g}, \bar{\pi})$ on M of the same type as (g, π) and with harmonic asymptotics (all in the same chart), such that we have*

$$\|g - \bar{g}\|_{W^{2,p}_{-q}} < \epsilon \quad \textit{and} \quad \|\pi - \bar{\pi}\|_{W^{1,p}_{-1-q}} < \epsilon,$$

the strict dominant energy condition

$$\bar{\mu} > |\bar{J}|_{\bar{g}}$$

holds, and

$$|E - \bar{E}| < \epsilon \quad \textit{and} \quad |P - \bar{P}| < \epsilon.$$

Remark. If we assume appropriate higher regularity for (M, g, π), then $(M, \bar{g}, \bar{\pi})$ will have the same regularity. This follows from applying Schauder estimates throughout the proof. Also, our proof shows that $(\bar{g}, \bar{\pi})$ can be chosen to be of type (p, q, q_0', α) for any given $q_0' > q_0$.

The proof of the density theorem consists of two general constructions. In Section 6.1 we deform (g, π) to an initial data set that satisfies a *strict* dominant energy inequality $\bar{\mu} > (1+\gamma)|\bar{J}|_{\bar{g}}$ for some constant $\gamma > 0$. In Section 6.2 we apply a cut-off argument as in [8, Theorem 1] to perturb to harmonic asymptotics, arguing that we can preserve the strict dominant energy condition in the process. Observe that even if we do not ask for strictness in the statement of Theorem 18, our proof still requires us to first perturb to strict dominant energy as in Section 6.1.

We first include a general fact that ADM energy and ADM linear momentum are continuous on the space of asymptotically flat initial data sets, as defined in Section 2.

Proposition 19 (*cf.* [18, Proposition 2.4])**.** *Let (g, π) and $(\bar{g}, \bar{\pi})$ be asymptotically flat initial data sets of type (p, q, q_0, α). Given $\epsilon > 0$ there exists $\delta > 0$ depending* *only on* $\epsilon, n, p, q, q_0, \|(g-\delta,\pi)\|_{W^{2,p}_{-q}\times W^{1,p}_{-q-1}}$, *and* $\|(\mu,J)-(\bar\mu,\bar J)\|_{W^{0,1}_{-n-q_0/2}}$ *such that if*

$$\|g-\bar g\|_{W^{2,p}_{-q}} \le \delta \quad \text{and} \quad \|\pi-\bar\pi\|_{W^{1,p}_{-1-q}} \le \delta,$$

then

$$|E-\bar E| < \epsilon \quad \text{and} \quad |P-\bar P| < \epsilon.$$

The argument is now standard and goes back to [26, p. 50] (for E only) and [8, p. 198] in the case of vacuum data. We include the proof for the sake of completeness. They key point is that continuity holds as long as we assume appropriate decay of the constraints.

Proof. By the definition of the energy E and the divergence theorem, we have that

$$\begin{aligned}
2(n-1)\omega_{n-1}E &= \lim_{r\to\infty}\int_{|x|=r}\sum_{i,j}(g_{ij,i}-g_{ii,j})\nu_0^j\,d\mathcal{H}_0^{n-1}\\
&= \int_{|x|=r}\sum_{i,j}(g_{ij,i}-g_{ii,j})\nu_0^j\,d\mathcal{H}_0^{n-1}\\
&\quad+\int_{|x|\ge r}\sum_{i,j}(g_{ij,ij}-g_{ii,jj})\,dx
\end{aligned}$$

for any r sufficiently large, and similarly for $\bar E$. Note that $\sum_{i,j}(g_{ij,ij}-g_{ii,jj}) = 2\mu + O(|x|^{-2-2q})$, since it differs from the scalar curvature R_g by terms quadratic in $\partial_k g_{ij}$ and π_{ij}. Since $(\mu-\bar\mu)\in W^{0,1}_{-n-q_0/2}$, it follows that there exists r_0 large and depending only on $\epsilon, n, p, q, q_0, \|\mu-\bar\mu\|_{W^{0,1}_{-n-q_0/2}}, \|(g-\delta,\pi)\|_{W^{2,p}_{-q}\times W^{1,p}_{-q-1}}$, and $\|(\bar g-\delta,\bar\pi)\|_{W^{2,p}_{-q}\times W^{1,p}_{-q-1}}$ such that for all $r\ge r_0$

$$\int_{|x|\ge r}\Big|\sum_{i,j}g_{ij,ij}-g_{ii,jj}-(\bar g_{ij,ij}-\bar g_{ii,jj})\Big|\,dx < (n-1)\omega_{n-1}\epsilon.$$

For the difference of the boundary integrals note that

$$\begin{aligned}
&\Big|\int_{|x|=r}\sum_{i,j}\left[g_{ij,i}-g_{ii,j}-(\bar g_{ij,i}-\bar g_{ii,j})\right]\nu_0^j\,d\mathcal{H}_0^{n-1}\Big|\\
&\le 2\sum_{i,j,k}\int_{|x|=r}|\partial_k(g-\bar g)_{ij}|\,d\mathcal{H}_0^{n-1}\\
&\le 2\mathcal{H}_0^{n-1}(\{|x|=r\})\sum_{i,j,k}\sup_{|x|=r}|\partial_k(g-\bar g)_{ij}|(x)\\
&\le 2(\omega_{n-1}r^{n-1})n^3(r^{-1-q}C_p\|g-\bar g\|_{W^{2,p}_{-q}}). \qquad (34)
\end{aligned}$$

where C_p is the constant governing the continuous embedding $W^{1,p}_{-1-q} \subset \mathcal{C}^{1-\frac{n}{p}}_{-1-q}$ for $p > n$ in Euclidean space. By choosing $\delta > \|g - \bar{g}\|_{W^{2,p}_{-q}}$ sufficiently small (depending on r) we can ensure that (34) is less than $(n-1)\omega_{n-1}\epsilon$ so that in total $|E - \bar{E}| < \epsilon$, as desired. The proof for the linear momentum is similar. □

6.1. **Perturbing to strict dominant energy condition.** We define the constraint map

$$\Phi(g, \pi) = (-2\mu, J) = \left(-R_g + |\pi|_g^2 - \tfrac{1}{n-1}(\operatorname{tr}_g \pi)^2, \operatorname{div}_g \pi\right).$$

It is also convenient to define the modified Lie derivative $\mathcal{L}_g Y = L_Y g - (\operatorname{div}_g Y) g$ of a vector field Y.

The proof of the following lemma is essentially the same as the proof of the surjectivity of $D\Phi|_{(g,\pi)}$ in [8, Proposition 3.1].

Lemma 20. *Suppose* $(g - \delta, \pi) \in W^{2,p}_{-q} \times W^{1,p}_{-1-q}$. *Define the linear map* $A : W^{2,p}_{-q} \times W^{1,p}_{-1-q} \to W^{0,p}_{-2-q}$ *by*

$$A(h, w) = D\Phi|_{(g,\pi)}(h, w) - (0, \tfrac{1}{2} h^l_j J_l),$$

where $D\Phi|_{(g,\pi)}$ *is the linearization of* Φ *at* (g, π). *Then* A *is surjective.*

Notation. For better readability, we define

$$s := \frac{4}{n-2},$$

which will appear frequently in this section.

For $(u - 1, Y) \in W^{2,p}_{-q}$, let

$$\tilde{g} = u^s g \quad \text{and} \quad \tilde{\pi} = u^{s/2}(\pi + \mathcal{L}_g Y).$$

Define the operator

$$T(u, Y) = (-2\tilde{\mu} u^s, \tilde{J} u^{s/2}),$$

where $(\tilde{\mu}, \tilde{J})$ are the mass and energy densities of $(\tilde{g}, \tilde{\pi})$. Explicitly, the components of T are given by

$$\begin{aligned} -2\tilde{\mu} u^s &= \tfrac{4(n-1)}{n-2} u^{-1} \Delta_g u - R_g - \tfrac{1}{n-1}\left(\operatorname{tr}_g \pi + \operatorname{tr}_g \mathcal{L}_g Y\right)^2 \\ &\quad + \left(|\pi|_g^2 + 2(\mathcal{L}_g Y)_{kl}\pi^{kl} + |\mathcal{L}_g Y|_g^2\right) \\ \tilde{J}_j u^{s/2} &= (\operatorname{div}_g \mathcal{L}_g Y + \operatorname{div}_g \pi)_j + (n-1)\frac{s}{2} u^{-1} u_{,k} (\pi + \mathcal{L}_g Y)^k_j \\ &\quad - \frac{s}{2} u^{-1} u_{,j} \operatorname{tr}_g(\pi + \mathcal{L}_g Y), \qquad j = 1, 2, \ldots, n. \end{aligned} \tag{35}$$

Here, all indices are raised with respect to g.

Lemma 21. *Suppose that* $(g-\delta,\pi) \in W^{2,p}_{-q} \times W^{1,p}_{-1-q}$. *For any* $(f,V) \in W^{0,p}_{-2-q}$ *there exist* $(v,Z) \in W^{2,p}_{-q}$ *and symmetric* $(0,2)$*-tensors* h, w *in* $\mathcal{C}^{3,\alpha}_c$ *so that*

$$DT|_{(1,0)}(v,Z) + D\Phi|_{(g,\pi)}(h,w) = (f, V_j + \tfrac{1}{2}h^l_j J_l).$$

If in addition $(f,V) \in \mathcal{C}^{0,\alpha}_{-n-q_0}$ *for some* $q_0 > 0$ *and* $\alpha > 0$, *and* $(g,\pi) \in \mathcal{C}^{2,\alpha}_{loc} \times \mathcal{C}^{1,\alpha}_{loc}$, *then we have* $(v,Z) \in \mathcal{C}^{2,\alpha}_{2-n}$.

Proof. The linearization of T at $(1,0)$ is

$$\begin{aligned} DT|_{(1,0)}(v,Z) = \Big(&\tfrac{4(n-1)}{n-2}\Delta_g v - 4Z_{k;l}\pi^{kl} - \tfrac{2}{n-1}\operatorname{tr}_g \pi \operatorname{div}_g Z, \\ &\operatorname{div}_g(\mathcal{L}_g Z)_j + \frac{(n-1)s}{2} v_{,k}\pi^k_j - \frac{s}{2} v_{,j} \operatorname{tr}_g \pi \Big) \end{aligned} \tag{36}$$

where all covariant derivatives and raising of indices are with respect to g. For $0 < a < (n-2)$ and $p > n$, $DT|_{(1,0)} : W^{2,p}_{-a} \to W^{0,p}_{-2-a}$ is Fredholm with index zero, *cf.* [5]. Because the linear map A defined in Lemma 20 is surjective onto $W^{0,p}_{-2-q}$ we can find $\mathcal{C}^{3,\alpha}$ compactly supported symmetric tensor fields $\{(h_k, w_k)\}_{k=1}^N$ whose images $A(h_k, w_k)$ span a subspace that complements the image of $DT|_{(1,0)}$ in $W^{0,p}_{-2-q}$. It follows that for every $(f,V) \in W^{0,p}_{-2-q}$ there exist $(v,Z) \in W^{2,p}_{-q}$ and $(h,w) \in \operatorname{span}\{(h_k,w_k)\}_{k=1}^N$ so that

$$DT|_{(1,0)}(v,Z) + D\Phi|_{(g,\pi)}(h,w) = (f, V_j + \tfrac{1}{2}h^l_j J_l). \tag{37}$$

This completes the first part of the proof.

Now suppose that $(f,V) \in \mathcal{C}^{0,\alpha}_{-n-q_0}$ for some $q_0 > 0$, and $(g,\pi) \in \mathcal{C}^{2,\alpha}_{loc} \times \mathcal{C}^{1,\alpha}_{loc}$. We view the system (37) as $(n+1)$ Poisson equations in v, Z_i with a decaying nonhomogeneous term. Note that the contribution $D\Phi|_{(g,\pi)}(h,w)$ to the nonhomogeneous term is a compactly supported Hölder function. Using [20, Theorem 2], which treats decay of solutions of the Poisson equation, and combining it with weighted Schauder estimates and a bootstrapping argument, we can conclude that $(v,Z) \in \mathcal{C}^{2,\alpha}_{2-n}$.

□

Theorem 22. *Let* (M^n, g, π) *be an asymptotically flat initial data set of type* (p, q, q_0, α), *as defined in Section 2. Assume that the dominant energy condition* $\mu \geq |J|_g$ *holds. For any* $\delta > 0$, *there exists asymptotically flat initial data* $(\bar{g}, \bar{\pi})$ *of the same type such that*

$$\|g - \bar{g}\|_{W^{2,p}_{-q}} \leq \delta \qquad \|\pi - \bar{\pi}\|_{W^{1,p}_{-1-q}} \leq \delta,$$

and for some $\gamma > 0$ *depending on* δ,

$$\bar{\mu} > (1+\gamma)|\bar{J}|_{\bar{g}}.$$

Proof. Choose a $\mathcal{C}^{3,\alpha}$ positive function f such that $f \equiv |x|^{-n-\min(1,q_0)}$ near infinity, and let $(v, Z) \in \mathcal{C}^{2,\alpha}_{2-n}$ and $(h, w) \in \mathcal{C}^{3,\alpha}_c$ be a solution of the system

$$DT|_{(1,0)}(v, Z) + D\Phi|_{(g,\pi)}(h, w) = (f, \tfrac{1}{2}h^l_j J_l),$$

whose existence is guaranteed by Lemma 21 (with $V \equiv 0$). Our goal is to show that for sufficiently small $t > 0$, the formula

$$\bar{g} = (1 + tv)^s(g + th) \quad \text{and} \quad \bar{\pi} = (1 + tv)^{s/2}(\pi + t\mathcal{L}_g Z + tw), \tag{38}$$

gives the desired initial data $(\bar{g}, \bar{\pi})$ in the statement of the theorem. Since we clearly have $\|g - \bar{g}\|_{W^{2,p}_{-q}} \le \delta$ and $\|\pi - \bar{\pi}\|_{W^{1,p}_{-1-q}} \le \delta$ for small enough t, it suffices to show that $\bar{\mu} > (1 + \gamma)|\bar{J}|_{\bar{g}}$ for some $\gamma > 0$ that depends on t.

In the following we denote $u = 1 + tv$ and define

$$\Phi_1(1 + tv, tZ, th, tw) = (-2\bar{\mu}u^s, \bar{J}u^{s/2}).$$

By Taylor expansion,

$$\begin{aligned}
&\Phi_1(1 + tv, tZ, th, tw)\\
&= \Phi_1(1, 0, 0, 0) + D\Phi_1|_{(1,0,0,0)}(v, Z, h, w) + \mathcal{R}\\
&= (-2\mu, J) + tDT|_{(1,0)}(v, Z) + tD\Phi|_{(g,\pi)}(h, w) + \mathcal{R}\\
&= (-2\mu, J) + t(f, \tfrac{1}{2}h^k_i J_k) + \mathcal{R},
\end{aligned} \tag{39}$$

where there is a minor abuse of notation in the last line, and the remainder term $\mathcal{R} = \mathcal{R}(x, t)$ has the form

$$\mathcal{R}(x, t) = t\int_0^1 \left[D\Phi_1|_{(1,0,0,0)+rt(v,Z,h,w)} - D\Phi_1|_{(1,0,0,0)}\right] (v, Z, h, w)\, dr.$$

Claim. *We have that*

$$|\mathcal{R}(x, t)| \le Ct^2(1 + |x|)^{2-2n} = O(t^2 f).$$

where C is a constant that does not depend on x or t.

Proof of claim. Clearly, $\mathcal{R}(x, t) = O(t^2)$, so it suffices to work outside a large ball B that contains the support of (h, w). Note that when $x \in M \smallsetminus B$ we have that $D\Phi_1|_{(1,0,0,0)+rt(v,Z,h,w)}(v, Z, h, w) = DT_{(1,0)+rt(v,Z)}(v, Z)$ and hence

$$\mathcal{R}(x, t) = t\int_0^1 \left[DT|_{(1,0)+rt(v,Z)} - DT|_{(1,0)}\right] (v, Z)\, dr.$$

From (35) we see that the linearization of T at (u, Y) is given by

$$\begin{aligned}
&DT|_{(u,Y)}(v, Z)\\
&= \Bigg(\tfrac{4(n-1)}{n-2}(-u^{-2}v\Delta_g u + u^{-1}\Delta_g v)\\
&\qquad - \tfrac{2}{n-1}(\operatorname{tr}_g \pi + \operatorname{tr}_g \mathcal{L}_g Y) \operatorname{tr}_g \mathcal{L}_g Z + 2(\pi + \mathcal{L}_g Y)^{kl}(\mathcal{L}_g Z)_{kl},\\
&\qquad \operatorname{div}_g(\mathcal{L}_g Z)_j + (n-1)\frac{s}{2}(vu^{-1})_{,k}(\pi + \mathcal{L}_g Y)^k_j + (n-1)\frac{s}{2}u^{-1}u_{,j}(\mathcal{L}_g Z)^k_j\\
&\qquad - \frac{s}{2}(vu^{-1})_{,j} \operatorname{tr}_g(\pi + \mathcal{L}_g Y) - \frac{s}{2}u^{-1}u_{,j} \operatorname{tr}_g \mathcal{L}_g Z\Bigg).
\end{aligned}$$

The claim follows from substituting $(1, 0) + rt(v, Z)$ and $(1, 0)$ for (u, Y), using that $(v, Z) \in \mathcal{C}^{2,\alpha}_{2-n}$, and estimating each term in an obvious way. ☐

From (39) we have

$$u^s\bar{\mu} = \mu + tf + O(t^2 f) \quad \text{and} \quad u^{s/2}\bar{J}_i = J_i + t\tfrac{1}{2}h^k_i J_k + O(t^2 f).$$

In particular, for sufficiently small $t > 0$, we have

$$u^s\bar{\mu} > \mu + \frac{t}{2}f \quad \text{everywhere on } M. \tag{40}$$

We claim that $u^s|\bar{J}|_{\bar{g}} < |J|_g + \frac{tf}{4}$ for $t > 0$ sufficiently small. Choose f_1 to be a $\mathcal{C}^{3,\alpha}$ nonnegative function whose support is larger than that of h. Then

$$\bar{g}^{ij} = u^{-s}(g^{ij} - tg^{ik}h^j_k + O(t^2 f_1))$$

Therefore

$$\begin{aligned}
(u^s|\bar{J}|_{\bar{g}})^2 &= u^{2s}\bar{g}^{ij}\bar{J}_i\bar{J}_j\\
&= (g^{ij} - tg^{ik}h^j_k + O(t^2 f_1))(J_i + t\tfrac{1}{2}h^k_i J_k + O(t^2 f))(J_j + t\tfrac{1}{2}h^k_j J_k + O(t^2 f))\\
&= g^{ij}J_iJ_j + t(-g^{ik}h^j_k J_iJ_j + g^{ij}\tfrac{1}{2}h^k_i J_kJ_j + g^{ij}J_i\tfrac{1}{2}h^k_j J_k)\\
&\quad + O(t^2|J|^2 f_1 + t^2|J|f + t^3 f_1 + t^4 f^2)\\
&= |J|^2_g + O(t^2|J|f + t^3 f^2)\\
&= \left(|J|_g + \frac{tf}{4}\right)^2 - \frac{tf}{2}|J|_g - \frac{t^2 f^2}{16} + O(t^2|J|f + t^3 f^2)\\
&< \left(|J|_g + \frac{tf}{4}\right)^2,
\end{aligned}$$

where we choose $t > 0$ to be sufficiently small in the last line, proving the claim. Observe that the computation above explains the motivation behind the definition of A. The $\frac{1}{2}h^l_j J_l$ term in A is chosen specifically so that the coefficient of t above, which is the first order change in $|J|_g$ under the deformation, vanishes.

Now fix $t > 0$ small enough so that $u^s\bar{\mu} > \mu + \frac{tf}{2}$ and $u^s|\bar{J}|_{\bar{g}} < |J|_g + \frac{tf}{4}$. Our assumptions imply that $\sup_M \frac{|J|_g}{f} < \infty$. For $x \in M$ such that $|\bar{J}|_{\bar{g}}(x) \neq 0$ it follows that

$$\frac{\bar{\mu}}{|\bar{J}|_{\bar{g}}} = \frac{u^s\bar{\mu}}{u^s|\bar{J}|_{\bar{g}}} > \frac{\mu + tf/2}{|J|_g + tf/4} \geq \frac{|J|_g + tf/2}{|J|_g + tf/4} \geq 1 + \gamma$$

where $\gamma^{-1} := 1 + \frac{4}{t}\sup_M \frac{|J|_g}{f}$. Since $\bar{\mu} > 0$ we conclude that $\bar{\mu} > (1+\gamma)|\bar{J}|_{\bar{g}}$ on M, as desired. Finally, note that γ depends only on t and $\sup_M \frac{|J|_g}{f}$. $\square$

6.2. **Harmonic asymptotics.** We first show that any asymptotically flat initial data set can be slightly perturbed so that

$$g = u^s\delta, \quad \pi = u^{s/2}\mathcal{L}_\delta Y,$$

outside a compact set, for some choice of u and Y, meanwhile prescribing any constraints (μ, J) that are close to the original ones. After that, we will show that if the imposed constraints decay rapidly enough, then (g, π) has harmonic asymptotics.

The first result is a modification of [8, Theorem 1], which treated the case $n = 3$, $\mu = \bar{\mu} = 0$, and $J = \bar{J} = 0$.

Lemma 23. *Suppose* $(g - \delta, \pi) \in W^{2,p}_{-q} \times W^{1,p}_{-1-q}$. *There exist* $C_0, \delta_0 > 0$ *so that given* $(\bar{\mu}, \bar{J}) \in W^{0,p}_{-2-q}$ *with* $\|(\bar{\mu} - \mu, \bar{J} - J)\|_{W^{0,p}_{-2-q}} \leq \delta \leq \delta_0$, *there is an initial data set* $(\bar{g}, \bar{\pi})$ *of the form*

$$g = u^s\delta, \quad \pi = u^{s/2}\mathcal{L}_\delta Y, \tag{41}$$

outside a compact set, for some $(u - 1, Y) \in W^{2,p}_{-q}$, *such that the mass and current densities of* $(\bar{g}, \bar{\pi})$ *are* $(\bar{\mu}, \bar{J})$. *Moreover,*

$$\|\bar{g} - g\|_{W^{2,p}_{-q}} \leq C_0\delta \qquad \|\bar{\pi} - \pi\|_{W^{1,p}_{-1-q}} \leq C_0\delta.$$

Proof. For $\lambda \geq 1$ large define the cut-off initial data

$$(\hat{g}_\lambda)_{ij} = \chi_\lambda g_{ij} + (1 - \chi_\lambda)\delta_{ij}, \qquad \hat{\pi}_\lambda = \chi_\lambda \pi,$$

where $\chi_\lambda(x) = \chi(x/\lambda)$ and χ is a $\mathcal{C}^{3,\alpha}$ cut-off function on $\mathbb{R}^n$ that is 1 on $\{|x| \leq 1\}$ and 0 on $\{|x| \geq 2\}$. Note that $\|(\hat{g}_\lambda - g, \hat{\pi}_\lambda - \pi)\|_{W^{2,p}_{-q} \times W^{1,p}_{-1-q}} \to 0$ as $\lambda \to \infty$.

In the following, we usually suppress the subscript λ when the context is clear. Define

$$\tilde{g} = u^s\hat{g} \qquad \tilde{\pi} = u^{s/2}(\hat{\pi} + \mathcal{L}_{\hat{g}}Y),$$

where $(u-1,Y) \in W^{2,p}_{-q}$. Let $(\tilde{\mu}, \tilde{J})$ be the mass and current densities of $(\tilde{g}, \tilde{\pi})$. Define the map $T_{(\hat{g},\hat{\pi})}(u,Y) = (-2\tilde{\mu}, \tilde{J})$ from $(W^{2,p}_{-q}+1)\times W^{2,p}_{-q} \to W^{0,p}_{-2-q}$. The linearization of $T_{(\hat{g},\hat{\pi})}$ at $(1,0)$ is

$$\begin{aligned}
&DT_{(\hat{g},\hat{\pi})}|_{(1,0)}(v,Z)\\
&= \Bigg(\tfrac{4(n-1)}{n-2}\Delta_{\hat{g}} v + s\big[R_{\hat{g}} - |\hat{\pi}|^2_{\hat{g}} + \tfrac{1}{n-1}(\operatorname{tr}_{\hat{g}}\hat{\pi})^2\big] v\\
&\qquad - 4Z_{k;l}\hat{\pi}^{kl} - \tfrac{2}{n-1}\operatorname{tr}_{\hat{g}}\hat{\pi}\,\operatorname{div}_{\hat{g}} Z,\\
&\qquad \operatorname{div}_{\hat{g}}(\mathcal{L}_{\hat{g}}Z)_j + (n-1)\frac{s}{2}v_{,k}\hat{\pi}^k_j - \frac{s}{2}v_{,j}\operatorname{tr}_{\hat{g}}\hat{\pi} - \frac{s}{2}(\operatorname{div}_{\hat{g}}\hat{\pi})_j v \Bigg),
\end{aligned}$$

where indices are raised and covariant derivatives are taken with respect to $\hat{g}$. The map $T_{(g,\pi)} : (W^{2,p}_{-q}+1)\times W^{2,p}_{-q} \to W^{0,p}_{-2-q}$ is defined analogously. Because $q \in (0, n-2)$ and $p > n$, $DT_{(\hat{g},\hat{\pi})}|_{(1,0)}$ and $DT_{(g,\pi)}|_{(1,0)}$ are Fredholm operators of index 0 for λ sufficiently large (see [5]).

Let K_1 be a complementing subspace for the kernel of $DT_{(g,\pi)}|_{(1,0)}$ in $W^{2,p}_{-q}\times W^{1,p}_{-2-q}$. Since the linearization $D\Phi|_{(g,\pi)} : W^{2,p}_{-q} \times W^{1,p}_{-1-q} \to W^{0,p}_{-2-q}$ is surjective (see the proof for $n=3$ in [8], which can be generalized to $n>3$) and because $DT_{(g,\pi)}|_{(1,0)}$ is Fredholm we can find $\mathcal{C}^{3,\alpha}$ compactly supported symmetric $(0,2)$-tensors $\{(h_k,w_k)\}_{k=1}^N$ whose images $D\Phi|_{(g,\pi)}(h_k,w_k)$ form a basis for a complementing subspace of the image of $DT_{(g,\pi)}|_{(1,0)}$ in $W^{0,p}_{-2-q}$. Let $K_2 = \operatorname{span}\{(h_k,w_k)\}_{k=1}^N$. We define the maps $\overline{T}_{(\hat{g},\hat{\pi})}, \overline{T}_{(g,\pi)} : K_1 \times K_2 \to W^{0,p}_{-2-q}$ by

$$\overline{T}_{(\hat{g},\hat{\pi})}(u,Y,h,w) = \Phi(u^s\hat{g} + h, u^{s/2}(\hat{\pi} + \mathcal{L}_{\hat{g}}Y) + w),$$

and

$$\overline{T}_{(g,\pi)}(u,Y,h,w) = \Phi(u^s g + h, u^{s/2}(\pi + \mathcal{L}_{g}Y) + w).$$

Note that the maps $\overline{T}_{(\hat{g},\hat{\pi})}, \overline{T}_{(g,\pi)}$ are continuously differentiable. Using that $(\hat{g},\hat{\pi})$ converges to (g,π) in $W^{2,p}_{-q}\times W^{1,p}_{-q-1}$ as $\lambda\to\infty$ it is easy to see that $D\overline{T}_{(\hat{g},\hat{\pi})}|_{(u,Y,h,w)}$ converges to $D\overline{T}_{(g,\pi)}|_{(u,Y,h,w)}$ as $\lambda\to\infty$ locally uniformly in $(u,Y,h,w)\in K_1\times K_2$ in the strong operator topology. Observe that $D\overline{T}_{(g,\pi)}|_{(1,0,0,0)}$ is an isomorphism by construction. We conclude from the inverse function theorem that there exists $\delta_0>0$ such that for all $\lambda\ge 1$ sufficiently large, $\overline{T}_{(g,\pi)}$ restricts to a $\mathcal{C}^1$ diffeomorphism defined on an open neighborhood of $(1,0,0,0)$ (independent of $\lambda \ge 1$) in $K_1\times K_2$ and onto an open neighborhood containing the $W^{0,p}_{-2-q}$-ball of radius $2\delta_0$ centered at $(-2\hat{\mu}, \hat{J})$.

Using that $\overline{T}_{(g,\pi)}(1,0,0,0) = (-2\mu, J)$ and that $\|(\mu,J)-(\hat{\mu},\hat{J})\|_{W^{0,p}_{-2-q}} \to 0$ as $\lambda\to\infty$ we see that $\|(\bar{\mu},\bar{J})-(\hat{\mu},\hat{J})\|_{W^{0,p}_{-2-q}} < 2\delta$ provided that $\|(\bar{\mu},\bar{J})-(\mu,J)\|_{W^{0,p}_{-2-q}} < \delta$ and $\lambda\ge 1$ is sufficiently large. Hence if $\delta\in(0,\delta_0)$ there exist $(u-1,Y)\in W^{2,p}_{-q}$ and

$\mathcal{C}^{3,\alpha}$ compactly supported symmetric $(0,2)$-tensors (h,w) such that $\overline{T}_{(\hat{g},\hat{\pi})}(u,Y,h,w) = (-2\bar{\mu},\bar{J})$ and such that

$$\|(u-1,Y)\|_{W^{2,p}_{-q}} \le C_1\delta \qquad \|(h,w)\|_{W^{2,p}_{-q}\times W^{1,p}_{-1-q}} \le C_1\delta,$$

for some constant $C_1 > 0$ that only depends on (g,π). The lemma follows easily from this.

□

The following proposition would be completely straightforward, but for a subtlety that arises in the case $n=3$.

Proposition 24. *Suppose that (M^n,g,π) is an asymptotically flat initial data set of type (p,q,q_0,α) such that*

$$g = u^s\delta, \quad \pi = u^{s/2}\mathcal{L}_\delta Y, \tag{42}$$

outside a large ball B for some $(u-1,Y)\in W^{2,p}_{-q}$. Furthermore, assume that $q_0>1$ (rather than simply $q_0>0$). Then (g,π) has harmonic asymptotics in the sense of Section 2.

Proof. By (35), outside a compact set, $(u-1,Y)$ satisfies

$$\begin{gathered}\tfrac{4(n-1)}{n-2}u^{-1}\Delta_\delta u - \tfrac{1}{n-1}\left(\operatorname{tr}_\delta \mathcal{L}_\delta Y\right)^2 + |\mathcal{L}_\delta Y|^2 = -2\mu u^s\\ \Delta_\delta Y_j + (n-1)\frac{s}{2}u^{-1}u_{,k}(\mathcal{L}_\delta Y)^k_j - \frac{s}{2}u^{-1}u_{,j}\operatorname{tr}_\delta(\mathcal{L}_\delta Y) = J_j u^{s/2},\end{gathered}$$

for $j=1,2,\ldots,n$. By Sobolev embedding, $(u-1,Y)=O^{1,\alpha}(|x|^{-q})$. Just as in the proof of Lemma 21, we may view this system as $n+1$ Poisson equations for the functions u, Y_i, so that $\Delta_\delta(u,Y) = O^\alpha(|x|^{\max(-2q-2,-n-q_0)})$. Then using the fact that $q>\frac{n-2}{2}$, Theorem 2 of [20] and weighted Schauder estimates imply that

$$\begin{aligned}u &= 1 + a|x|^{2-n} + O^{2,\alpha}(|x|^{2-n-\gamma})\\ Y_i &= b_i|x|^{2-n} + O^{2,\alpha}(|x|^{2-n-\gamma})\end{aligned} \tag{43}$$

for constants a, b_i, for any $0<\gamma<\min(2q+2-n,q_0)$, outside B. In particular $(u-1,Y)\in\mathcal{C}^{2,\alpha}_{2-n}$ outside B. For $n>3$, repeating the above argument using $(u-1,Y)=O^{1,\alpha}(|x|^{2-n})$ yields the desired result, since $\max(2-2n,-n-q_0)<-n-1$. For $n=3$ we argue as follows.

We expand the nonlinear "source" terms above. We find that

$$\begin{aligned}|\mathcal{L}_\delta Y|^2 &= |L_Y g|^2 - 2\operatorname{tr}_\delta(L_g Y)(\operatorname{div}_\delta Y) + 3(\operatorname{div}_\delta Y)^2\\ &= 2\left[\sum_{i,j=1}^3 Y_{i,j}^2 + Y_{i,j}Y_{j,i}\right] - (\operatorname{div}_\delta Y)^2\end{aligned}$$

while

$$(\mathrm{tr}_\delta\,\mathcal{L}_\delta Y)^2 = (\mathrm{div}_\delta\, Y)^2.$$

Since

$$Y_{i,j} = -b_i \frac{x_j}{|x|^3} + O^{1,\alpha}(|x|^{-2-\gamma}),$$

we have

$$(\mathrm{div}_\delta\, Y)^2 = \left(\sum_{i=1}^{3} Y_{i,i}\right)^2 = \frac{(b\cdot x)^2}{|x|^6} + O^{1,\alpha}(|x|^{-4-\gamma})$$

and

$$\sum_{i,j=1}^{3} Y_{i,j}^2 + Y_{i,j}Y_{j,i} = \frac{|b|^2}{|x|^4} + \frac{(b\cdot x)^2}{|x|^6} + O^{1,\alpha}(|x|^{-4-\gamma}).$$

Thus the constraint equation for μ becomes

$$\Delta_\delta u = -\frac{2|b|^2}{|x|^4} - \frac{3(b\cdot x)^2}{2|x|^6} + O^{\alpha}(|x|^{-4-\gamma}),$$

as long as $\gamma < q_0$. Now observe that

$$\Delta_\delta\left[\frac{x_i x_j}{|x|^4}\right] = \frac{2\delta_{ij}}{|x|^4} - \frac{4x_i x_j}{|x|^6} \qquad \text{for } i,j\in\{1,2,3\},$$
$$\Delta_\delta\left[\frac{1}{|x|^2}\right] = \frac{2}{|x|^4}.$$

For appropriate choice of constants C_0 and C_{ij}, we have

$$\Delta_\delta\left[u - C_0\frac{1}{|x|^2} - C_{ij}\frac{x_i x_j}{|x|^4}\right] = O^{\alpha}(|x|^{-4-\gamma}).$$

Now apply [20, Theorem 2] and weighted Schauder estimates to obtain the desired result for u. The argument for Y is analogous. $\square$

6.3. **Proof of Theorem 18.** Using Theorem 22 and Proposition 19 we may reduce to the case where $\mu > (1+\gamma)|J|_g$. For $l \geq 1$ define $\xi_l(x) := \xi(\frac{x}{l})$ where

$$\xi(x) = \begin{cases} 1 & \text{when } |x| \leq 1 \\ |x|^{-1} & \text{when } |x| \geq 2 \end{cases}.$$

Note that $\|\xi_l(\mu,J) - (\mu,J)\|_{\mathcal{C}^{0,\alpha}_{-n-q_0}}$ and hence $\|\xi_l(\mu,J) - (\mu,J)\|_{W^{0,p}_{-2-q}}$ tends to zero as $l\to\infty$. Also note that $\xi_l(\mu,J)\in\mathcal{C}^{0,\alpha}_{-n-q_0-1}$, so that it has improved decay. By Lemma 23 and Proposition 24 there exist initial data sets (g_l,π_l) with harmonic asymptotics whose mass and current densities are $\xi_l(\mu, J)$, and such that $\|(g-g_l, \pi-\pi_l)\|_{W^{2,p}_{-q}\times W^{1,p}_{-1-q}} \to 0$ as $l \to \infty$. In particular, $\|g-g_l\|_{C^0} \to 0$ so that

$$|\xi_l J|^2_{g_l} = g_l^{ij}\xi_l^2 J_i J_j = \xi_l^2 |J|^2_g (1+o(1)) \quad \text{as} \quad l \to \infty,$$

and hence

$$\xi_l \mu > \xi_l(1+\gamma)|J|_g \geq \left(1+\frac{\gamma}{2}\right)|\xi_l J|_{g_l},$$

for l sufficiently large. This means that the strict dominant energy condition holds for (g_l, π_l). The convergence of mass and linear momentum as $l \to \infty$ follows from Proposition 19. $\square$

Remark. One subtlety in the proof of Theorem 18 above is that the decay conditions for (μ, J) improve in the step where we are reducing to harmonic asymptotics. In fact, if we are content to preserve the dominant energy condition without requiring strictness, we can arrange for $(\bar{\mu}, \bar{J})$ to vanish outside a large compact set by choosing ξ with compact support.

Remark. Finally we remark that it is possible to prove the positive mass theorem from the positive energy theorem. We argue by contradiction. Assuming the dominant energy condition and $E < |P|$, we can use the previous remark and [8] to approximate our initial data set by one which is identical with a suitably chosen slice in a Kerr spacetime outside a compact set, such that we still have the dominant energy condition and $E < |P|$. This makes it possible to further deform the data in the Kerr region to a "standard" slice with negative ADM energy, contradicting the positive energy theorem. Note that the positive energy theorem was proven in [27] for $n = 3$ and the proof generalized to the cases $4 \leq n \leq 7$ in [11].

References

1. William K. Allard, *On the first variation of a varifold: boundary behavior*, Ann. of Math. (2) **101** (1975), 418–446. MR 0397520 (53 #1379)
2. Lars Andersson, Marc Mars, and Walter Simon, *Stability of marginally outer trapped surfaces and existence of marginally outer trapped tubes*, Adv. Theor. Math. Phys. **12** (2008), no. 4, 853–888. MR 2420905 (2010a:83005)
3. Lars Andersson and Jan Metzger, *The area of horizons and the trapped region*, Comm. Math. Phys. **290** (2009), no. 3, 941–972. MR 2525646 (2010f:53118)
4. Abhay Ashtekar and Gregory J. Galloway, *Some uniqueness results for dynamical horizons*, Adv. Theor. Math. Phys. **9** (2005), no. 1, 1–30. MR 2193368 (2006k:83101)
5. Robert Bartnik, *The mass of an asymptotically flat manifold*, Comm. Pure Appl. Math. **39** (1986), no. 5, 661–693. MR 849427 (88b:58144)
6. Robert Beig and Piotr T. Chruściel, *Killing vectors in asymptotically flat space-times. I. Asymptotically translational Killing vectors and the rigid positive energy theorem*, J. Math. Phys. **37** (1996), no. 4, 1939–1961. MR 1380882 (97d:83033)

7. Piotr T. Chruściel and Daniel Maerten, *Killing vectors in asymptotically flat space-times. II. Asymptotically translational Killing vectors and the rigid positive energy theorem in higher dimensions*, J. Math. Phys. **47** (2006), no. 2, 022502, 10. MR 2208148 (2007b:83054)
8. Justin Corvino and Richard M. Schoen, *On the asymptotics for the vacuum Einstein constraint equations*, J. Differential Geom. **73** (2006), no. 2, 185–217. MR 2225517 (2007e:58044)
9. Lu Ding, *Positive mass theorems for higher dimensional Lorentzian manifolds*, J. Math. Phys. **49** (2008), no. 2, 022504, 12. MR 2392853 (2008m:53081)
10. Frank Duzaar and Klaus Steffen, *λ minimizing currents*, Manuscripta Math. **80** (1993), no. 4, 403–447. MR 1243155 (95f:49062)
11. Michael Eichmair, *The Jang equation reduction in dimensions less than eight*, preprint.
12. ———, *The Plateau problem for marginally outer trapped surfaces*, J. Differential Geom. **83** (2009), no. 3, 551–583. MR 2581357
13. ———, *Existence, regularity, and properties of generalized apparent horizons*, Comm. Math. Phys. **294** (2010), no. 3, 745–760. MR 2585986 (2011d:53171)
14. Michael Eichmair and Jan Metzger, *Jenkins-Serrin-Spruck type results for Jang's equation*, preprint (2011).
15. Gregory J. Galloway, *Rigidity of marginally trapped surfaces and the topology of black holes*, Comm. Anal. Geom. **16** (2008), no. 1, 217–229. MR 2411473 (2009e:53087)
16. Gregory J. Galloway and Niall Ó Murchadha, *Some remarks on the size of bodies and black holes*, Classical Quantum Gravity **25** (2008), no. 10, 105009, 9. MR 2416045 (2009c:83058)
17. Gregory J. Galloway and Richard Schoen, *A generalization of Hawking's black hole topology theorem to higher dimensions*, Comm. Math. Phys. **266** (2006), no. 2, 571–576. MR 2238889 (2007i:53078)
18. Lan-Hsuan Huang, *On the center of mass in general relativity*, arXiv:1101.0456 (2011).
19. Joachim Lohkamp, *The higher dimensional positive mass theorem I*, arXiv:math/0608795v1 (2006).
20. Norman Meyers, *An expansion about infinity for solutions of linear elliptic equations.*, J. Math. Mech. **12** (1963), 247–264. MR 0149072 (26 #6568)
21. Richard Schoen, *Uniqueness, symmetry, and embeddedness of minimal surfaces*, J. Differential Geom. **18** (1983), no. 4, 791–809 (1984). MR 730928 (85f:53011)
22. ———, *Variational theory for the total scalar curvature functional for Riemannian metrics and related topics*, Topics in calculus of variations (Montecatini Terme, 1987), Lecture Notes in Math., vol. 1365, Springer, Berlin, 1989, pp. 120–154. MR 994021 (90g:58023)
23. ———, *Mean curvature in Riemannian geometry and general relativity*, Global theory of minimal surfaces, Clay Math. Proc., vol. 2, Amer. Math. Soc., Providence, RI, 2005, pp. 113–136. MR 2167257 (2006f:53044)
24. Richard Schoen and Leon Simon, *A new proof of the regularity theorem for rectifiable currents which minimize parametric elliptic functionals*, Indiana Univ. Math. J. **31** (1982), no. 3, 415–434. MR 652826 (84j:49039)
25. Richard Schoen and Shing Tung Yau, *On the proof of the positive mass conjecture in general relativity*, Comm. Math. Phys. **65** (1979), no. 1, 45–76. MR 526976 (80j:83024)
26. ———, *The energy and the linear momentum of space-times in general relativity*, Comm. Math. Phys. **79** (1981), no. 1, 47–51. MR 609227 (82j:83045)
27. ———, *Proof of the positive mass theorem. II*, Comm. Math. Phys. **79** (1981), no. 2, 231–260. MR 612249 (83i:83045)

28. Richard M. Schoen and Shing Tung Yau, *Complete manifolds with nonnegative scalar curvature and the positive action conjecture in general relativity*, Proc. Nat. Acad. Sci. U.S.A. **76** (1979), no. 3, 1024–1025. MR 524327 (80k:58034)
29. Leon Simon, *Lectures on geometric measure theory*, Proceedings of the Centre for Mathematical Analysis, Australian National University, vol. 3, Australian National University Centre for Mathematical Analysis, Canberra, 1983. MR 756417 (87a:49001)
30. Jean E. Taylor, *The structure of singularities in solutions to ellipsoidal variational problems with constraints in* R^3, Ann. of Math. (2) **103** (1976), no. 3, 541–546. MR 0428182 (55 #1208b)
31. Brian White, *The space of m-dimensional surfaces that are stationary for a parametric elliptic functional*, Indiana Univ. Math. J. **36** (1987), no. 3, 567–602. MR 905611 (88k:58027)
32. ———, *Existence of smooth embedded surfaces of prescribed genus that minimize parametric even elliptic functionals on* 3*-manifolds*, J. Differential Geom. **33** (1991), no. 2, 413–443. MR 1094464 (92e:58048)
33. Edward Witten, *A new proof of the positive energy theorem*, Comm. Math. Phys. **80** (1981), no. 3, 381–402. MR 626707 (83e:83035)
34. P. F. Yip, *A strictly-positive mass theorem*, Comm. Math. Phys. **108** (1987), no. 4, 653–665. MR 877642 (88m:83029)

Massachusetts Institute of Technology, eichmair@math.mit.edu

Columbia University, lhhuang@math.columbia.edu

Queens College and CUNY Graduate Center, dan.lee@qc.cuny.edu

Stanford University, schoen@math.stanford.edu